

Biodiversity of Larsemann Hills, Antarctica

Gupta P^{*}

Scientist, Ecology Section, Botanical Survey of India, Ministry of Environment Forests & Climate Change, Government of India, CNH Building, Botanic Garden, Howrah 711 103, India

^{*}**Correspondence to :** Ecology Section, Botanical Survey of India, Ministry of Environment Forests & Climate Change, Government of India, CNH Building, Botanic Garden, Howrah 711 103, India, e-mail: drpratibha2011@rediffmail.com

Publication History

Received: 09 April 2015

Accepted: 18 May 2015

Published: 1 July 2015

Citation

Gupta P. Biodiversity of Larsemann Hills, Antarctica. *Climate Change*, 2015, 1(3), 174-183

ABSTRACT

Antarctica is the coldest, driest place and fifth-largest continent on this Earth and contains many unique geological, glaciological and biological features. The Larsemann Hills (69°20'S to 69°30'S Latitude., 75°55'E to 76°30'E Longitude) is located approximately halfway between Vest fold Hills and Amery Ice Shelf on South-eastern coast of Prydz Bay. Human activities in Larsemann Hills is promoted due to its coastal location, ice free landscape, Australian summer research base (Law base), Chinese research station (Zhongshan) and Russian research stations (Progress) were established within the area of 3 km from each other on eastern broknes. After that there was rapid infrastructure development in the area and further scientific research and the potential for tourist visits resulted in notable localised alteration of the environment, ultimately affect the biodiversity. The Antarctic environment / climate is highly susceptible to the impacts of human activities and has much less natural ability to recover from disturbance than the environment of other continents. These changes ultimately affect the environment and growth of organisms. Larsemann Hills represent the southernmost coastal oasis contains diverse flora and fauna. In flora - Mosses, Lichens and Algae were observed. Among algae mainly Cosmarium, Diatoms, Dinoflagellates and Cyanobacterial mats were observed from different water bodies and terrestrial habitats. As for as fauna is concerned breeding sea bird like Snow petrels, Wilson's Storm Petrel, South polar Skuas were found. Besides this, Seals, Adelie Penguin and Emperor Penguin were occasionally observed in Larsemann Hills area. However, very little is known about the terrestrial micro fauna. In lakes and streams species of protozoans, platyhelminths, rotifers, tardigrades, nematodes, arthropods, etc. have been reported.

Keywords: Biodiversity, Larsemann Hills, Antarctica

1. INTRODUCTION

Antarctica is almost completely covered in ice and it is a gift of nature in natural forms. This untouched continent attracts scientific workers to study the nature in natural form. There are two seasons summer and winter. 1st December to 28th February is summer and 1st March to 30th November is winter period. On the basis of duration of light the day is of six months and night is of six month. There is no permanent human

Gupta,
Biodiversity of Larsemann Hills, Antarctica,
Climate Change, 2015, 1(3), 174-183,
www.discovery.org.in

population and however, various Antarctic research stations from many nations that are inhabited by summer and winters teams of scientists with Leaders on rotation for scientific work.

2. LARSEMANN HILLS

The Larsemann Hills is an ice-free area located approximately halfway between Vestfold Hills and Amery Ice Shelf on South-eastern coast of Prydz Bay (Fig 1 & 2). This area has a low, gentle and rolling topography merging with the polar ice cap in the south-southeast and surrounded by sea in other three directions. It is punctuated with small islands to the north and northeast.




Figure 1
Map of Larsemann Hills, Antarctica

The region consists of two main peninsulas (Stornes and Broknes), together with other peninsulas and a number of scattered offshore islands. At 40 km², the Larsemann Hills is the second largest of only four major ice free oasis. The highest elevations are around 180 m above sea level (ANARE, 2000). The hills are dissected by steep valleys lying between the ice sheet and the coast. Many valley floors are filled with multi-year ice but those that are ice-free contain a complex geomorphological history preserved in erosional features, including glacial striae and tafoni, glacial and glacio fluvial sediments (Stuwe et al., 1989). More than 150 freshwater lakes are found in the Larsemann hills (Gillieson et al., 1990) ranging from small ephemeral ponds to large water bodies such as Progress Lake (10 ha and 3.8 m deep). Some of these water bodies are briefly ice-free or partially ice-free in the summer months when their temperatures increase rapidly; surface water in some of the shallower ones reaches more than 8°C. For the remainder of the year (8 - 10 months) they are covered with about 2 m of ice. Some lakes have evidence of past shorelines which are up to 2 m above present levels but do not exhibit the highly elevated salinities associated with lakes in the Rauer Islands (Hodgson et al., 2001). Lake systems are typically closed and contained in steep-sided V-shaped valleys normally around 50 - 100 m deep and less than 1 km long. These valleys dissect the area and provide a conduit for summer melt water streams between the few lake systems that are open. During Antarctic summer in December, January and February the daily air temperature frequently exceeds 4°C and has been known to reach 10°C (ANARE 2000). Mean monthly winter temperature is between -15°C and -18°C. Precipitation occurs as snow and is unlikely to exceed 250 mm water equivalent annually. Strong, katabatic winds blow most mornings. Preliminary studies by McMinn and Hodgson (unpublished) and Ellis-Evans et al. (1998) have revealed that lakes in the Larsemann Hills contain a great diversity of biological and

physical markers from which past environments can be inferred. Some of the most scientifically important lakes on eastern Broknes and other lakes of Larsemann Hills area are collectively recognised as the area of most important ecological feature. These lakes are particularly valuable for their relatively simple natural ecosystems and are susceptible to physical, chemical and biological modification within their catchable boundaries.



Figure 2
 Satellite Imagery Map of Larsemann Hills, Antarctica

Coastal location and ice free landscape in Larsemann Hills area attracted human activity as tourist visit and its potentials for scientific researches induced human activity is affecting the diversity of this area. The Antarctic climate/environment is super susceptible to the impacts of human activities and has negligible natural ability to recover from disturbance than the environment of other continents. So the primary study of biodiversity in this area before impact of human activity is of valuable significance.

Microclimatic conditions and presence of freshwater lakes during summer provide hospitable environment able to support Antarctic life forms. Biodiversity from diverse habitats of Larsemann Hills have been studied by some workers of different countries. However, the work by Indian workers is still in very nascent stage as Indian Research Station Bharati was recently established in March, 2012.

3. DIVERSITY OF FLORA AND FAUNA

Larsemann Hills coastal oasis contains diverse flora and fauna. In flora - Mosses, Lichens, Algae and Bacteria were observed. In algae mainly Cosmarium, Diatoms, Dinoflagellates and Cyanobacterial mats were observed from different water bodies and terrestrial habitats. As for as fauna is concerned, breeding sea bird like *Pagodroma nivea* (Snow Petrels), *Oceanites oceanicus* (Wilson's Storm Petrel), *Catharacta maccormicki* (South Polar Skuas) was found. Besides this, *Leptonychotes weddelli* (Weddell Seals), *Pygoscelis adeliae* (Adelie Penguin) and *Aptenodytes forsteri* (Emperor Penguin) were occasionally observed in Larsemann Hills area. However, very little is known about the terrestrial micro fauna. In lakes and streams species of Protozoans, Platyhelminths, Rotifers, Tardigrades, Nematodes, Arthropods, etc. have been reported.

As per the report, the Larsemann Hills indicate the flora of the Ingrid Christensen Coast is relatively uniform and restricted to a distribution of bryophytes, lichens and terrestrial algae. It is believe that the nature of basement rock, the relatively recent exposure from the ice cap and prevailing wind direction in greater Prydz Bay area contribute to the less than 1 % of the Larsemann Hills has vegetative cover.

Terrestrial life forms including bryophytes, lichens and accompanying invertebrates are found inland from the coast. Nevertheless large moss beds are known to occur in sheltered sites on the larger island associated with *Pygoscelis adeliae* (Adelie penguin) moulting sites and nunataks in the southwest.

4. BRYOPHYTES

Bryophytes are represented by widely spread terrestrial mosses in a mosaic of habitats like soils, outside the snow fields, marginal zones or bank of the water bodies and melted water streams, sheltered habitats of exposed knobs with representative but low to moderate snow accumulation, biogenic remains and around the nests of *Catharacta maccormicki* (south polar Skua). Bryophytes also found in liquid phase of water and among the running water they become biological indicator of the boundary of water. The presence of mosses on nunataks and in mountain ranges depends on the availability of water. The presence of few species of mosses in such type of environment indicates their ecological limits. Altogether seven species of mosses and one species of liverwort (ASAM 6, 2007) have been reported depicted in Table 1.

Table 1

Bryo-diversity of Larsemann Hills, Antarctica

S. No.	Name of the Species	Family	Type
1	<i>Bryum algens</i> Cardot	Bryaceae	Moss
2	<i>Bryum argenteum</i> Hedw.	Bryaceae	Moss
3	<i>Bryum pseudotriquetrum</i> (Hedw.) Schwaegr	Bryaceae	Moss
4	<i>Ceratodon purpureus</i> (Hedw.) Brid.	Ditrichaceae	Moss
5	<i>Grimmia antarctici</i> Card	Grimmiaceae	Moss
6	<i>Grimmia lawiana</i> J.H. Willis	Grimmiaceae	Moss
7	<i>Sarconeurum glaciale</i> (C.Bryhn.) Card. Et Bryhn	Pottiaceae	Moss
8	<i>Cephaloziella exiliflora</i> H.Streimann	Cephaloziellaceae	Liverwort

Out of seven species of mosses, *Bryum pseudotriquetrum* (Hedw.) Schwaegr is found most abundantly. However, only one species of liverwort *Cephaloziella exiliflora* H.Streimann is reported on the unnamed outcrop south of Stornes Peninsula and known from only four other Antarctic localities.

5. LICHENS

Lichens species have been reported from diverse habitats of McLeod Island of Larsemann Hills, Antarctica. Eighteen species of lichens from McLeod Island was earlier studied by Australian researchers. Rod Seppelt studies the lichens of whole Ingrid Christensen Coast, Prydz Bay and listed 25 species of lichens including four species (*Acarospora gwynnii* C.W. Dodge & E.D. Rudolph, *Buellia frigida* Darb., *Candelariella flava* (C.W. Dodge & G.E. Baker) Castello & Nimis and *Umbilicaria decussate* (Vill.) Zahlbr.) from Larsemann Hills which was remain unpublished. After that, Singh et al., 2007 reported 25 lichen species from this area. Recently, lichen flora of this region studied by Rai et al., 2012 and comprises 27 species (Table 2).

Table 2

Lichen diversity from McLeod Island of Larsemann Hills, Antarctica

S. No.	Name of the Species	Family	Growth	Substratum
1	<i>Acarospora gwynnii</i> C.W. Dodge & E.D. Rudolph	Acarosporaceae	Crustose	Rock
2	<i>Arthonia lapidicola</i> (Taylor) Branth & Rostr.	Arthoniaceae	Crustose	Rock
3	<i>Buellia frigida</i> Darb.	Caliciaceae	Crustose	Rock
4	<i>Buellia grimmiae</i> Filson	Caliciaceae	Crustose	Soil, Moss
5	<i>Caloplaca athallina</i> Darb.	Teloschistaceae	Crustose	Moss
6	<i>Caloplaca citrina</i> (Hoffm.) Th. Fr.	Teloschistaceae	Crustose	Rock
7	<i>Caloplaca lewis-smithii</i> Søchting & Øvstedal	Teloschistaceae	Crustose	Moss

8	<i>Caloplaca saxicola</i> (Hoffm.) Nordin	Teloschistaceae	Crustose	Rock, Soil
9	<i>Candelariella flava</i> (C.W. Dodge & G.E. Baker) Castello & Nimis	Candelariaceae	Crustose	Rock, Soil, Moss
10	<i>Carbonea vorticosa</i> (Flörke) Hertel	Lecanoraceae	Crustose	Rock
11	<i>Huea coralligera</i> (Hue) C.W. Dodge & G.E. Baker	Teloschistaceae	Crustose	Moss
12	<i>Lecanora expectans</i> Darb.	Lecanoraceae	Crustose	Soil, Moss
13	<i>Lecanora geophila</i> (Th. Fr.) Poelt	Lecanoraceae	Crustose	Moss
14	<i>Lecidea cancriformis</i> C.W. Dodge & G.E. Baker	Lecideaceae	Crustose	Rock, Soil
15	<i>Lecidella patavina</i> (A. Massal.) Knoph & Leuckert	Lecanoraceae	Crustose	Rock, Soil
16	<i>Lecidella siplei</i> (C.W. Dodge & G.E. Baker) May. Inoue	Lecanoraceae	Crustose	Rock
17	<i>Physcia caesia</i> (Hoffm.) Hampe ex Fűrnr.	Physciaceae	Foliose	Rock, Moss
18	<i>Physcia dubia</i> (Hoffm.) Lettau	Physciaceae	Foliose	Rock
19	<i>Pseudephebe minuscula</i> (Nyl. Ex Arnold) Brodo & D. Hawksw.	Parmeliaceae	Fruticose	Rock, Soil
20	<i>Rhizoplaca melanophthalma</i> (Ram.) Leuckert and Poelt	Lecanoraceae	Crustose	Rock, Soil, Moss
21	<i>Rinodina olivaceobrunnea</i> C.W. Dodge & G.E. Baker	Physciaceae	Crustose	Soil, Moss
22	<i>Rinodina peloleuca</i> (Nyl.) Müll. Arg.	Physciaceae	Crustose	Rock
23	<i>Sarcogyne privigna</i> (Ach.) A. Massal.	Acarosporaceae	Crustose	Rock, Soil
24	<i>Umbilicaria decussata</i> (Vill.) Zahlbr.	Umbilicariaceae	Foliose	Rock
25	<i>Usnea antarctica</i> Du Rietz	Parmeliaceae	Fruticose	Rock
26	<i>Xanthoria elegans</i> (Link) Th. Fr.	Teloschistaceae	Foliose	Rock
27	<i>Xanthoria mawsonii</i> C.W. Dodge	Teloschistaceae	Foliose	Rock

Crustose lichens were dominant with 20 species. *Physcia caesia* (Hoffm.) Hampe ex Fűrnr., *Physcia dubia* (Hoffm.) Lettau, *Xanthoria elegans* (Link) Th. Fr. and *Xanthoria mawsonii* C.W. Dodge were the only foliose lichen in this area. However, only two species viz. *Pseudephebe minuscula* (Nyl. ex Arnold) Brodo & D. Hawksw. and *Usnea antarctica* Du Rietz. have been reported as fruticose lichen.

6. ALGAE

Australia, United Kingdom (Hodgson et al., 2001) Canada (Hannington, 2002) Belgium (Sabbe et al., 2003), etc. are working on algal flora on various Antarctic regions and presented comparative accounts of algal diversity with their respective countries and also in different part of Antarctic region. But very little work has been done on studies on algal diversity from diverse habitats of Larsemann hills, Antarctica.

Algae occurs even in wide range of habitats and have been distributed all over land and water system often in such an environments where there is no other vegetation possibly due to their adaptive capability to extreme adverse environmental conditions with respect to different climatic / environmental factors, availability of nutrients, etc. Their occurrence even in wide range of ecologically stress conditions and extreme habitats proves that they are very tolerant. It occurs in fresh-water ecosystem like lakes, ponds, rivers, wetland, etc. and marine water system

like salt marshes and pans, estuaries, brackish waters and ocean. Besides, it also occurs on stones, rocks, snow and in cold lakes, and on objects which remain moisten and get solar light from any angle even for a short span of time.

Diatoms, from fresh and saline water lakes of Larsemann Hills and Rauer Island have been studied (Sabbe et al. 2003 and Sarah, 2010) (Table 3).

Table 3

Algal diversity of Larsemann Hills, Antarctica

S. No.	Name of the Species	Class	Family
1	<i>Diademsis costei</i> Le Cohu & Van de Vijver	Bacillariophyceae	Diadesmidaceae
2	<i>Diademsis langebertalotii</i> Le Cohu & Van de Vijver	Bacillariophyceae	Diadesmidaceae
3	<i>Diademsis subantarctica</i> Le Cohu & Van de Vijver	Bacillariophyceae	Diadesmidaceae
4	<i>Diatomella balfouriana</i> Greville	Bacillariophyceae	Pinnulariaceae
5	<i>Psammothidium marginulatum</i> (Grunow) L. Bukhtiyarova & Round	Bacillariophyceae	Achnanthesiaceae
6	<i>Gomphonema parvulum</i> (Kütz.) Kützing	Bacillariophyceae	Gomphonemataceae
7	<i>Planothidium delicatum</i> (Kütz.) Round & Bukhtiyarova	Bacillariophyceae	Achnanthesiaceae

Phytoplanktons mostly comprises autotrophic nanoflagellates, dinoflagellates, desmids belongs to genus *Cosmarium* occurs in lakes. Study reveals that heterotrophic nanoflagellates are more common in comparison to autotrophic nanoflagellates exhibiting low species diversity. Only three or four species in most lakes are particularly abundant in shallow lakes. *Parphysomonas* is very common in shallow lakes. Ciliates like *Strombidium* are most common genera and found in less quantity. Species of *Holyophyra* found in most lakes of this area. *Coscinodiscus*, *Epithemia*, *Nitzschia*, *Thalassiosira*, *Oscillatoria* and *Chaetoceros* have been reported from Larsemann Hills, Antarctica. *Coscinodiscus* and *Nitzschia* were most abundant and widely distributed respectively. Extensive cyanobacterial mats was observed in most of the lakes which has accumulated since ice retreat and is consequently thicker on the islands and thinner in young lakes adjacent to the polar plateau. These cyanobacterial (blue-green) mats are of exceptional thickness not generally observed in other freshwater system of Antarctica. Cyanobacterial mats are also widely distributed in streams and wet seepage area.

7. BACTERIA

The Antarctic continent shows microbial community development because of its special environment, geographic isolation and little anthropogenic influence. The Culture-independent community analysis reveals bacterial diversity. The dominant taxa in the glacier forefields are Actinobacteria, Acidobacteria, Proteobacteria, Bacteroidetes, and Chloroflexi. The connection of soil characteristics with bacterial community structure showed that soil parameter and soil formation along the glacier forefield influence the distribution of certain phyla (Bajerski and Wagner, 2013). Several groups like Proteobacteria or Gemmatimonadetes depend on water availability, whereas the most dominant group of the forefields the Actinobacteria is related to the presence of trace elements. Two new psychrotolerant bacteria, *Herbaspirillum psychrotolerans* and *Chryseobacterium frigidisoli* were found as novel species of the family of Oxalobacteraceae and Flavobacteriaceae, respectively. They are able to grow at low temperatures tolerating temperature fluctuations and not specialised to a certain substrate and well-adapted to the cold and oligotrophic environment. The prokaryotic community of Antarctic soil is estimated as diverse and variable (Barrett et al., 2006, Niederberger et al., 2008). The dominant groups are Bacteroidetes, Actinobacteria, Proteobacteria, *Deinococcus/Thermus*, Acidobacteria, Firmicutes and Cyanobacteria (Aislabie et al., 2006). *Herbaspirillum psychrotolerans* sp. nov., a member of the family Oxalobacteraceae from a glacier forefield (Bajerski, 2013b). *Chryseobacterium frigidisoli* sp. nov., a psychrotolerant species of the family Flavobacteriaceae isolated from sandy permafrost from a glacier forefield (Bajerski, 2013a).

8. SEA BIRDS

Ecological and biological studies at Broknes on the south polar skua i. e. *Catharacta maccormicki* have been carried out (Wang, 1991). Study indicates that Larsemann Hills area also provide breeding sites for other seabirds. Breeding sea bird like Snow petrels (*Pagodroma nivea*), Wilson's Storm Petrels (*Oceanites oceanicus*), south polar Skuas (*Catharacta maccormicki*) and Weddell seals haul out close to sea shore to breed and moult (Wang, 1991). These three species of seabird breed within the Larsemann Hills. Location of breeding pairs and numbers are recorded from eastern Broknes but their distribution is throughout the remaining area is not certain. Snow petrels (*Pagodroma nivea*), Wilson's Storm Petrels (*Oceanites oceanicus*) nests found from October to February in sheltered bedrock fragments, cervices, boulder slopes and rock

falls on broknes. South polar Skuas (*Catharacta maccormicki*) are present between mid-late October and early April nesting on broknes. Besides this, Adelie Penguin (*Pygoscelis adeliae*) and Emperor Penguin (*Aptenodytes forsteri*) were occasionally observed in Larsemann Hills area but no breeding colonies are found in this area. However, the birds visit from colonies of nearby islands during summer season to moult.

9. SEALS

Weddell seals (*Leptonychotes weddelli*) are numerous on Larsemann Hills coast using the sea ice in the area to pup from October and to moult from late December until March. Weddell seals hauled out in Thala Fjored and on rafted ice immediately to west of Stornes and numerous smaller groups amongst offshore islands and ice to north-east of broknes. Crabeater seals (*Lobodon carcinophagus*) and Leopard seals (*Hydrurga leptonyx*) are also occasional visitors of this area.

10. TERRESTRIAL MICRO FAUNA

Little work is done on the terrestrial micro fauna of Larsemann Hills. Lakes and streams provide many habitats which contains rich and varied fauna typical of Antarctic regions. Species of rotifer (including *Monogononta* and *Bedelloidea*), tardigrades, arthropods, protozoans, platyhelminths and nematodes have been reported (Dartnall, 1995). Rotifers occur sporadically in number of lakes. The cladoceran *Daphniopsis studeri* is one of few species of freshwater crustacean known to occur in the lakes of continental Antarctica has been identified in most Larsemann Hills lakes. Benthic species of deep water lakes of this area are dominated by thick cyanobacterial mats (Ellis-Evan et. al., 1998).

11. PROTOZOA

Protozoans have been reported from Scandent, Discussion, Sibthorpe, Progress, Reid and four unnamed lakes. Generally, seen on the bottom of the freshwater lakes were oval elliptical, transparent or pale brown ciliates, small ciliates with spikey tufts - holotrichia, large solitary paramecium-type ciliates and a few small amoebae.

12. PLATYHELMINTHES

Solitary specimen of small blind acoelomates was found in the sediments of lake Scandrett and Unnamed lake.

13. TARDIGRADA

Two species of smooth tardigrada were recorded from the freshwater lakes. These would appear to belong to the *Isohysibius* group reported from Discussion, Sibthorpe, Progress, Reid, and two unnamed lakes of Larsemann Hills. In addition *Hfilnesium tardigradum* Doyere was found in terrestrial mosses at the edge of lake Scandrett. Five genera of terrestrial tardigrade such as *Hysibius*, *Minibiotus*, *Diphascion*, *Milnesium* and *Pseudechiniscus* are reported in association with vegetation.

14. ROTIFERS

Seventeen species of rotifers viz. *Cephalodella sterea*, *Cephalodella ventripes*, *Collotheca ornate cornuta*, *Encentrum mustela*, *Encentrum spatium*, *Epiphanes senta*, *Lepadella patella*, *Lepadella acuminata*, *Notholca* sp., *Ptygura crystalline*, *Resticula gelida*, *Adineta grandis*, *Adineta* sp., *Habrotricha constricta* *Philodina gregaria* and two species of *Philodina* sp. have been reported from different lakes of Larsemann Hills, Antarctica.

15. NEMATODES

Terschellingia, *Araeolaimus*, *Axonolaimus*, *Chromadorella*, *Daptonema*, *Halalaimus*, *Paralinhomoeus*, *Sabatieria*, *Stiphonolaimus*, *Sphaerolaimus* and *Theristus* found as dominant nematode genera from near shore Antarctic locations in the Larsemann Ice Shelf (Ingole and Singh, 2010).

16. ARTHROPODA

Two species of arthropoda found in the freshwater lakes i.e. cladoceran *Daphniopsis studeri* Ruch and the copepod *Acanthocyclops mirnyi* Borutzky and Vinogradov. *Daphniopsis studeri* was the largest and most abundant species found. Large numbers of males were found in lake Scandrett during the February. This species is commonly recorded from the subantarctic islands of the Southern Indian Ocean. *Acanthocyclops mirnyi* was also found in the larger fresh water lakes. It was first found in the Bunger Hills (Korotkevich, 1958). Study of biodiversity of Larsemann Hills area is very important because of the rapidly increasing anthropogenic activity may bring about changes in diversity of flora and fauna and may also influence on micro and macro climatic conditions of this area.

ACKNOWLEDGEMENTS

The author is grateful to Dr. P. Singh, Director, Botanical Survey of India for encouragement and Director, NCAOR, Ministry of Earth Sciences, Goa for support and help during 33rd Indian Scientific expedition to Antarctica. I acknowledge the Satellite Image of Larsemann Hills with courtesy Google Earth.

REFERENCES

1. Aislabie JM, Chhour KL, Saul DJ, Miyauchi S, Ayton J, Paetzold RF, Balks MR. Dominant bacteria in soils of Marble Point and Wright Valley, Victoria Land, Antarctica. *Soil Biol Biochem*, 2006, 38, 3041-3056
2. ANARE. Visitor's guide to the Larsemann Hills (<http://www.antdiv.gov.au>), 2000
3. ASMA. Management Plan for Antarctic Specially Managed Area No. 6 Larsemann Hills, East Antarctica. *Measure 2 Annex B*, 2007
4. Bajerski F, Wagner D. Bacterial succession in Antarctic soils of two glacier forefields on Larsemann Hills, East Antarctica. *FEMS Microbial Ecology*, 2013, 85, 128-42
5. Bajerski F, Ganzert L, Mangelsdorf K, Padur L, Lipski A, Wagner D. *Chryseobacterium frigidisoli* sp. nov., a psychrotolerant species of the family Flavobacteriaceae isolated from sandy permafrost from a glacier forefield. *Int. J Syst Evol Microbiol*, 2013a, 63, 2666-71
6. Bajerski F, Ganzert L, Mangelsdorf K, Lipski A, Busse HJ, Padur L, Wagner D. *Herbaspirillum psychrotolerans* sp. nov., a member of the family Oxalobacteraceae from a glacier forefield. *Int. J Syst Evol Microbiol*, 2013b, 63, 3197-3203.
7. Barrett JE, Virginia RA, Hopkins DW, Aislabie J, Bargagli R, Bockheim JG, Campbell IB, Lyons WB, Moorhead DL. Terrestrial ecosystem processes of Victoria Land, Antarctica. *Soil Biol Biochem* 2006, 38, 3019-3034.
8. Dartnall HJG. Rotifers and other aquatic invertebrates from the Larsemann Hills, Antarctica. *Pap & Proc Roy Soc Tasmania*, 1995, 129, 17-23
9. Ellis-Evanjs C, Laybourn P, Ayliss YB, Perriss J. Physical, chemical and microbial community characteristics of lakes of the Larsemann Hills, Continental Antarctica. *Arch Hydrobiol*, 1998, 141, 209-230
10. Gillieson D, Burgesjs S, Spatea, Cochran. An atlas of the lakes of the Larsemann Hills, Princess Elizabeth Land, Antarctica. *ANARE Research Notes*, No 74, 1990, pp. 1-173
11. Hannington M. Joint German-Canadian Research on Seafloor Hydrothermal Activity in Antarctica: Sonne Cruise SO-155: 14, *News Letter for the Canadian Antarctic Research Network*, 2002, 14, 1-16
12. Hodgson DA, Noon PE, Vyverman W, Bryant CL, Gore DB, Appleby P, Gilmour M, Verleyen E, Sabbe K, Jones VJ, Ellis-Evans JC, Wood PB. Were the Larsemann Hills ice-free through the Last Glacial Maximum? *Antarc. Sci.* 2001, 13, pp. 440-454
13. Ingole B, Singh R. Biodiversity and community structure of free living marine nematodes from the Larsemann Ice Shelf, East Antarctica. *Curr Sci*, 2010, 99, 1413- 1419
14. Korotkevich YS. Nasaliniye vodocmovoazisov v estochnoy Antarktide. Inf. Byull. Sov. Antarkt. Eksped 3: 91 - 98 (Translated into English, 1964: Concerning the population of water bodies in the oasis of East Antarctica. *Sov Antarc Exped Inf Bull*, 1959, 1, 154-161
15. Niederberger TD, McDonald IR, Hacker AL, Soo RM, Barrett JE, Wall DH, Cary SC. Microbial community composition in soils of Northern Victoria Land, Antarctica. *Environ Microbiol*, 2008, 10, 1713-1724
16. Rai H, Khare R, Nayak S, Upreti DK, Gupta RK. Lichen Synusiae in East Antarctica (Schirmacher Oasis and Larsemann Hills): substratum and morphological preferences. *Czech Polar Reports* 2012, 1, 65-77
17. Sabbe K, Verleyen E, Hodgson DA, Vanhoutte K, Vyverman W. Benthic diatom flora of freshwater and saline lakes in the Larsemann Hills and Rauer Islands, East Antarctica. *Antarc Sci*, 2003, 15, 227-248
18. Sarah AS, De Vuver BV, Dominic A, Hodgson M. D. M, Verleyen E, Lee S. Diatoms as indicators of environmental change in Antarctic and subantarctic freshwaters. In *The Diatoms: Applications for the Environmental and Earth Sciences*, 2nd Edition, (eds. Sonol, JP. & Stroermer, EF.) Cambridge University Press. 2010, 14, 267-283
19. Singh SM, Nayaka S, Upreti DK. Lichen Communities in Larsemann Hills, East Antarctica. *Curr Sci*, 2007, 93, 1670-1672
20. Stuwiek BM, Peer H. Geology and structure of the Larsemann Hills area, Prydz Bay, East Antarctica. *Australian J Earth Sci*, 1989, 36, 2, 19-24.
21. Wang Z. Ecology of Catharacta maccormicki near Zhongshan Station in Larsemann Hills, East Antarctica. *Antarc Res (Chinese edition)*, 1991, 3, 45-55.



Indian Research Station Bharati, Larsemann Hills, Antarctica



Bryophytes from Larsemann Hills, Antarctica



Bryophytes from Larsemann Hills, Antarctica



Lichens from Larsemann Hills, Antarctica



Algae in water logged area and from water body of Larsemann Hills, Antarctica



Catharacta maccormicki (South polar Skua)



Pagodroma nivea (Snow petrel)



Oceanites oceanicus (Wilson's Storm Petrel)



Leptonychotes weddelli (Weddell Seal)



Pygoscelis adeliae (Adelie penguin)



Aptenodytes forsteri (Emperor Penguin)