



OPEN

High-resolution ecosystem changes pacing the millennial climate variability at the Middle to Upper Palaeolithic transition in NE-Italy

Federica Badino^{1,2✉}, Roberta Pini², Cesare Ravazzi², Milan Chytrý³, Paolo Bertuetti², Eugenio Bortolini¹, Lydie Dudová⁴, Marco Peresani^{2,5}, Matteo Romandini¹ & Stefano Benazzi¹

Observation of high-resolution terrestrial palaeoecological series can decipher relationships between past climatic transitions, their effects on ecosystems and wildfire cyclicality. Here we present a new radiocarbon dated record from Lake Fimon (NE-Italy) covering the 60–27 ka interval. Palynological, charcoal fragments and sediment lithology analysis were carried out at centennial to sub-centennial resolutions. Identification of the best modern analogues for MIS 3 ecosystems further enabled to thoroughly reconstruct structural changes in the vegetation through time. This series also represents an “off-site” reference record for chronologically well-constrained Palaeolithic sites documenting Neanderthal and *Homo sapiens* occupations within the same region. Neanderthals lived in a mosaic of grasslands and woodlands, composed of a mixture of boreal and broad-leaved temperate trees analogous to those of the modern Central-Eastern Europe, the Southern Urals and central-southern Siberia. Dry and other grassland types expanded steadily from 44 to 43 ka and peaked between 42 and 39 ka, i.e., about the same time when Sapiens reached this region. This vegetation, which finds very few reliable modern analogues in the adopted Eurasian calibration set, led to the expansion of ecosystems able to sustain large herds of herbivores. During 39–27 ka, the landscape was covered by steppe, desert-steppe and open dry boreal forests similar to those of the modern Altai-Sayan region. Both Neanderthal and Sapiens lived in contexts of expanded fire-prone ecosystems modulated by the high-frequency climatic cycles of MIS 3.

Marine isotope stage (MIS) 3 (~ 60–30 ka), a period of intermediate global ice volume between MIS 4 and the Last Glacial Maximum (~ MIS 2), was one of the periods of most unstable climate, closely interwoven with human evolution history. During MIS 3, climate variability was associated with abrupt atmospheric shifts over Greenland (Dansgaard–Oeschger [D–O] events), episodes of massive iceberg discharge and freshwater inputs into the North Atlantic up to the Iberian margin (Heinrich events [HEs])¹, which generated climatic and ecological responses worldwide (Heinrich stadials [HS])². Information about regional palaeoclimate during MIS 3 comes from isotope records of speleothems (e.g., Hölloch Cave, N-Alps³; Ascunsa Cave, S-Carpathians⁴, Pozzo Cucù Cave, Apulia, Southern Italy)⁵.

In Southern-Europe, along the Italian Peninsula (Monticchio, Castiglione, and Lagaccione, Italy)^{6–8} and in Mediterranean lowlands (Tenaghi Philippon and Megali Limni, Greece)^{9–11}, temperate forests and forest-steppe were supported during Greenland Interstadials (GI), with steppe expansion during Greenland Stadials (GS), further exacerbated during HS^{12,13}. Less extensive forest patches occurred in mountain areas during GSs (e.g., Ioannina, western Greece)¹⁴, where temperate tree population survival was favoured by orographic precipitation throughout MIS 3. Forests were also present in some parts of central and eastern Europe, e.g., in river valleys, especially in the Carpathians¹⁵. In more continental areas, such as the lower Danube Plain and the semi-arid

¹Department of Cultural Heritage, University of Bologna, 48121 Ravenna, Italy. ²CNR-Institute of Environmental Geology and Geoengineering, Lab. of Palynology and Palaeoecology, Research Group on Vegetation, Climate and Human Stratigraphy, 20126 Milan, Italy. ³Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic. ⁴Department of Paleocology, Institute of Botany, Czech Academy of Sciences, Brno, Czech Republic. ⁵Department of Humanities, Prehistoric and Anthropology Sciences, University of Ferrara, 44100 Ferrara, Italy. ✉email: federica.badino@igag.cnr.it

Pannonian Basin, vegetation formations became more open, likely due to drought stress¹⁶, and loess deposits formed repeatedly during MIS 3 (e.g., Nussloch and Willendorf sites)¹⁷. In western-central and western Europe, tundra vegetation prevailed during MIS 3 with relatively small stadial/interstadial fluctuations of tree and shrub populations: e.g., La Grande Pile, Les Echets, Bergsee (after 45 ka cal BP) and Füramoos pollen records¹⁸. Within this context, the replacement of Neanderthals by *Homo sapiens* is recorded across Europe in a diachronous and culturally articulated succession of local technocomplexes (e.g., Uluzzian), that replaced the Mousterian material cultures during the so called Middle to Upper Palaeolithic transition roughly between 45 and 40 ka cal BP¹⁹.

Relationships between climate fluctuations, environmental changes and the arrival of Sapiens at the expense of Neanderthals in Europe represent a hotly debated topic^{5,20}.

Several factors contribute to the persisting uncertainty on the timing and mode of the earliest migration of Sapiens in Europe and the environmental context where this happened. In fact, most of the currently available information are inferred from chronological correlation with geographically remote records, i.e., Greenland ice cores²¹, rather than from regional environmental series. Moreover, only a subset of the several European archaeological sites recording the Middle to Upper Palaeolithic transition have been accurately investigated, or are well-documented and precisely dated^{22,23}.

In the S–E fringe of the Italian Alps, in the sub-alpine area of the Berici Hills (S–E) and Lessini Mountains (N–W) (ca. 50 ka apart, Fig. 1b), favourable circumstances have fostered the accumulation and preservation of deposits documenting Neanderthals and Sapiens occupations at Fumane and Broion caves and shelters in the 45–40 ka time frame^{24–29}. In the same area, the natural archive of Lake Fimon (Fig. 1) provides a palaeoecological series that entirely covers the last Glacial-Interglacial cycle^{30–32} (Fig. 2). Due to its favourable position and stratigraphic continuity, this sequence represents a suitable “off-site”³³ reference record for NE-Italian Palaeolithic sites. In this paper, we present a new set of chronostratigraphic and high-resolution geochemical and microbotanical analyses carried out on the Fimon PD and TdA cores for the 60–27 ka interval (Figs. 1c and 2 and “Methods”). Specific objectives of this study are: (1) the reconstruction of millennial to sub-millennial environmental and fire regime dynamics throughout MIS 3, (2) the identification of the best modern analogues for MIS 3 ecosystems, and (3) the environmental contextualization of human occupation during the Middle to Upper Palaeolithic Transition in NE-Italy.




Figure 1. Geographical framework. (a) Location of high-resolution palaeoecological series entirely covering MIS 3 (yellow dots), speleothem series (light blue dots), loess profiles (light green dots) and Palaeolithic sites documenting the Uluzzian and Neronian cultures (red dots): (1) Fumane Cave; (2) Broion shelter; (3) Crvena Stijena; (4) Grotta del Cavallo; (5) Grotta di Uluzzo C; (6) Castelcivita; (7) Grotta della Cala; (8) Colle Rotondo; (9) Grotta La Fabbrica; (10) Klissoura Cave; (11) Kephalaria Cave; (12) Mandrin Caves. Sea level has lowered by 120 m during the LGM³⁴ and by 74 m during MIS 3 relative to the current sea level³⁵ with new estimates suggesting higher sea levels between -18 and -40 m asl during MIS 3³⁶. (b) The study area in NE Italy (Veneto region). Location of Lake Fimon and nearby Palaeolithic sites documenting the Uluzzian culture: Fumane cave (Lessini Mountains) and Broion shelter (Berici Hills), are highlighted: (c) Detail of the Lake Fimon area and of the coring sites (white dots): Ponte sulla Debba (PD), Torri di Arcugnano (TdA) and Fimon Lago (FL). Maps were created using ESRI ArcGIS 10.7 software (<https://www.esri.com/software/arcgis>) and the web page: <https://www.freeworldmaps.net/europe/italy/veneto.html>.




Figure 2. Summary of the palynostratigraphic data from Lake Fimon. **(a)** Synthetic composite pollen record obtained from cores FL (12–27 ka) and PD (> 27 ka) documenting the history of plant communities throughout the Late Pleistocene (modified after^{30,37}). **(b)** New high-resolution synthetic palaeoecological record from the Fimon PD core for the period 60–27 ka cal BP: conifers (green), broad-leaved trees (orange), upland herbs (light yellow), xerophytes (yellow).

Results

Chronology. The Fimon PD record was age-depth modelled between 60.1 and 26.7 ka cal BP (mean ages) over a 2.36 m long core section (i.e., 21.75–19.39 m). We relied on ^{14}C ages from two Lake Fimon sequences: Fimon PD and Fimon TdA cores, previously correlated for the Last Glacial-Interglacial cycle interval³⁷. In this study, a targeted correlation for the MIS 3 interval was carried out relying on high-resolution litho-, chrono- and biostratigraphic data, allowing a precise arrangement of all the available Fimon TdA ^{14}C dates along the Fimon PD record (see “Methods” section and Supplementary Fig. S1 for further details). The list of radiocarbon ages and the tie-point included in the model (Supplementary Fig. S3) is shown in Table 1. Deposits span ca. 33,300 years entirely covering Marine Isotope Stage 3. Chronological boundaries between pollen zones were identified at (95% probability): 43,096–41,079 cal BP (FPD1-2), 41,233–37,695 cal BP (FPD2-3), 33,885–30,098 cal BP (FPD3-4). The median resolution is 140 years for pollen and 80 years for sieved charcoal samples. Median accumulation rates are variable along the record: 0.006 cm/years for silty clay, gyttja and carbonate mud deposits (FL3-4) and 0.02 cm/years for peat interval (FL5), see Supplementary SI-1 and Supplementary Fig. S4 for further details.

Gradients in pollen composition and dominant vegetation formations. The Fimon PD fossil pollen record was analysed with a median time resolution of 140 years. Cluster analysis identified four pollen zones named FPD 1 to 4, the latter previously further subdivided into FPD 2 to 7³². Relatively high woody-pollen percentages (mean = 55%), dominated by *Pinus sylvestris/mugo* with peaks up to 93%, characterize the whole sequence (Fig. 2). Two major drops in forest cover occur in FPD2c and FPD4a pollen zones (Fig. 2). In the long term, land cover developed from a mosaic of grasslands and forest patches (FPD1), composed of a mixture of boreal and broad-leaved temperate trees, to dry boreal forests (FPD4) throughout more open phases dominated by herbs and xerophytes (FPD2-3). A notable change occurred at ca. 39 ka, i.e., the boundary between cluster groups FPD1-2 and FPD3-4, with samples showing positive and negative PCA1 scores, respectively (Figs. 2, 3).

The oldest pollen zone FPD1 is dominated by grasslands with patches of forests, including thermophilous *Ulmus* and *Quercus robur/cerris* type (up to 3%; Fig. 2), which expanded alongside *Picea*, whose percentages varied between 0.4 and 8% (Fig. 2). Among other temperate tree taxa, *Tilia* persisted up to ca. 39 ka, while *Alnus glutinosa* type, *Corylus* and *Betula* are documented throughout the whole record, the latter showing rapid increases in relationship with enhanced fire frequency (Fig. 2 and Supplementary Fig. S7). Pollen zone FPD2 is mainly characterized by an expansion of Cichorioideae (up to 31%) and other Asteraceae (e.g., *Anthemis* type, *Aster* type, *Xeranthemum inapertum* type, Supplementary Fig. S3). Pollen zones FPD 3–4 are characterized by fluctuations between boreal forests with *Pinus sylvestris/mugo* and *Larix*, the latter with a continuous presence since 31 ka, and open environments dominated by *Artemisia* (mean = 8%) and Chenopodiaceae (mean = 1.5%; Fig. 2).

Modern analogues. We used the MAT—Modern Analogue Technique to find the best-matching modern analogues for the Fimon PD fossil pollen assemblages. The adopted Eurasian calibration set consisted of 5978 modern pollen spectra: 5925 are from the Eurasian Modern Pollen Database vers. 2³⁹ and 53 from the Southern Urals, collected in 2007 (see “Methods” for details). We also used climate-sensitive pollen indicators to better describe past floristic structures and ecoclimatic gradients (see “Methods” section).

Overall, 83% of the closest (first) modern analogues are considered “good analogues”, i.e., sq. chord dist. < 5% (Supplementary Fig. S9, Supplementary Table S1)⁴⁰. Modern analogues for fossil spectra included in pollen zone FPD1 (60–42 ka) belong to the “Central European mixed forests” (29%), the “Sarmatic mixed forests” (16%) and the “Kazakh forest steppe” (14%). Such environments are characterized by open mixed forests with a contribution of temperate broadleaved trees such as *Quercus*, *Tilia* and *Ulmus*, and a mosaic of grasslands with patches of

Radiocarbon ages from Lake Fimon—Ponte sulla debba (PD) and Torri di Arcugnano (TdA) cores								
Lab code	Core	Lithology	Original depth (m)	Fimon PD depth (m)	Material dated	^{14}C Age BP	2 σ calibration range (cal years BP) IntCal20	Median probability (cal years BP)
UBA-7831	Fimon PD	Peat	19.40–19.41	19.40–19.41	Bulk sediment	22,593 ± 115	26,816–27,211	26,953
UBA-7830	Fimon PD	Peat	19.51–19.52	19.51–19.52	Bulk sediment	23,165 ± 116	27,249–27,686	27,439
UBA-7829	Fimon PD	Peat	19.74–19.755	19.74–19.755	Bulk sediment	24,376 ± 187	28,014–29,056	28,591
UBA-15493	Fimon TdA	Peat	26.95–26.96	19.85–19.86	Bulk sediment	26,158 ± 97	30,125–30,768	30,376
UBA-46815	Fimon TdA	Slightly organic gyttja	27.66–27.68	20.37–20.39	Charcoal	33,990 ± 532	37,306–40,225	38,898
UBA-46816	Fimon TdA	Silty clay	28.61–28.63	20.84–20.86	Charcoal and wood	37,523 ± 837	40,950–42,744	41,952
Biostratigraphic events from Lake Fimon—Ponte sulla debba (PD) record								
Biostratigraphic event	Core		Original depth (m)	Fimon PD depth (m)	Isotopic correlated event		Age cal ka and error	
<i>Picea</i> — <i>Tilia</i> rise	Fimon PD	Silty clay	21.72–21.75	21.72–21.75	Start of GI 17		60.2 ± 0.4 ^{30,38}	

Table 1. List of the radiocarbon ages and biostratigraphic events available from the Fimon PD and TdA sequences.




Figure 3. PCA ordination of terrestrial pollen taxa (>2%) and sites showing changes in the pollen patterns in terrestrial ecosystems. Photos from the Southern Urals (FPD1 and FPD2) and the Altai Mountains (FPD3 and FPD4) show landscapes analogous to NE-Italy around Lake Fimon during the four time periods (pollen zones FPD1 to 4).

Pinus sylvestris and deciduous trees. Other analogues belong to the “Sayan montane conifer forests” (20%) and the “Altai alpine meadow and tundra” (6%) ecoregions in semi-dry, mesic, wet and saline grasslands environments with both alpine and steppe species, and open larch forests. Species occurring frequently in the analogous Altai-Sayan grasslands and also found in NE Italy today include *Anthoxanthum alpinum*, *Aster alpinus*, *Bistorta major*, *B. vivipara*, *Galium verum* and *Rumex alpestris*, the moss *Rhytidium rugosum* and the lichen *Cetraria islandica*.

The 42–39 ka interval (pollen zone FPD2) is characterized by the marked expansion of Cichorioideae together with other Asteraceae (*Anthemis*, *Aster* type, *Xeranthemum inapertum* type), Ranunculaceae, *Geranium molle* type, *Ephedra fragilis* and *E. distachya* types and *Galium* (Supplementary Fig. S3). Such taxa, when combined, highlight a trend towards more open and dry conditions (see discussion). 48% of the pollen spectra from this pollen zone lack of good modern analogues; among the few statistically reliable modern analogues, several come from fallow lands, arid meadows and pasture habitats (Supplementary Table S1).

The modern analogues for the 39–31 ka interval (pollen zone FPD3) mostly belong to the “Altai alpine meadow and tundra” (39%), the “Central European mixed forests” (12%), the “Western Siberian hemiboreal forests” (12%), the “East European forest-steppe” (6%) and the “Western European broadleaf forests” (6%) ecoregions (Fig. 4). The best modern analogues in the Altai-Sayan region are again represented by a mosaic of various types of grasslands with woodland patches and tundra, the latter mainly at elevations of 2000–2200 m⁴¹. The Fimon PD fossil data are lacking or poorly representing tundra pollen indicators (e.g., *Dryas*, Ericaceae, *Salix*), suggesting a very little contribution of this habitat at Lake Fimon during MIS 3. Compared to the previous phases, dry steppe and desert-steppe sensu⁴² spread during pollen zone FPD 3 as documented by high *Artemisia* and *Chenopodiaceae* % values. Modern analogues include several species also found in northern Italy, e.g., *Allium strictum*, *Festuca valesiaca*, *Galium boreale* and *G. verum*, *Phleum phleoides*, *Stipa capillata* and *S. pennata*. Other analogues belong to taiga with *Larix*, *Picea* and *Pinus sibirica* (= *P. cembra* s. lat.), however, the latter species was always below 5% of total pollen in fossil spectra from Lake Fimon.

The youngest interval, i.e., 31–27 ka (pollen zone FPD4), finds its analogues especially in the “Altai alpine meadow and tundra” (62%), with a little contribution from the “East Siberian taiga” (18%) and the “Alps conifer and mixed forests” (8%) ecoregions (Fig. 4). The best modern analogues correspond to the same grassland, steppe and desert-steppe already detected for pollen zone FPD3, but several best analogues also come from forests, partly from *Larix-Picea-Pinus sibirica* taiga with *Vaccinium vitis-idaea* and abundant mosses, and partly from hemiboreal forests with *Pinus sylvestris*, *P. sibirica*, *Betula pendula* and a species-rich herb layer.

Fire frequency, return interval, and severity reconstruction. We investigated changes in fire activity through the record of sieved charcoal particles (i.e., 62–125 μm , 125–500 μm and > 500 μm fractions) with




Figure 4. (a) Location of the four nearest modern analogues selected for each pollen zone: pollen zone FPD1 (60–42 ka cal BP); FPD2 (42–39 ka cal BP); FPD3 (39–31 ka cal BP); FPD4 (31–27 ka cal BP), plotted on Biomes map (RESOLVE Ecoregions 2017)⁴³. (b) Eco-regions map⁴³ shows the regions selected by the first modern analogue for each pollen zone. (c) Pie-charts showing the proportion of eco-regions selected by the first analogue for each pollen zone. The grey sections represent values <5%. Maps were created using ESRI ArcGIS 10.7 software (<https://www.esri.com/software/arcgis>).

a median time resolution of 80 years. Charcoal Accumulation Rates (CHAR) of particles > 125 μm at depths between 21.75 and 19.39 m were calculated (Supplementary Fig. S7). Signal to noise index (SNI) is greater than the critical value of 3 which indicates suitability for peak analysis⁴⁴ for the 45% of the record (median = 2.7, min = 0 and max = 334.4; Supplementary Fig. S8). We avoided interpreting peaks corresponding to SNI < 3, mostly occurring in the lowermost part of the record (60–44 ka). Peak analysis revealed twenty-four statistically significant CHAR peaks showing variable magnitude between 0.0001 and 0.24 pieces cm⁻² peak⁻¹, with highest values between 31 and 27 ka. Fire return intervals range between 270 and 430 years and fire frequency up to 3 fires per 1000 years during phases of major biomass availability (GI 12, 8, 4 and 3 forest stages). No or low fire activity was registered during phases of prolonged biomass decrease and expansion of open vegetation (Supplementary Fig. S7).

Discussion

Vegetation dynamics and modern analogues for MIS 3 environments in NE-Italy. The best modern analogues for Fimon PD pollen spectra are mainly from the Eurasian temperate zone, especially from the transition zone between the Siberian boreal zone and the drylands of Central Asia. In particular, the Altai-Sayan area appears to be the best modern analogue ecoregion for most of the MIS 3 environments documented in the Fimon PD fossil record. This similarity has been already outlined for the Last Glacial ecosystems in Central Europe^{15,45}.

Although the landscape around Fimon was a mosaic of meadows, steppe and open forests throughout MIS 3, it changed over time, particularly in the proportion and composition of forest patches and the ratio of wet/mesic to dry grasslands. During the 60–42 ka interval (FPD1), several modern analogues suggest that the landscape may have been similar to the modern Eastern European forest-steppe such as that in the Southern Urals, where temperate broadleaved trees (*Quercus*, *Tilia*, *Ulmus*) are mixed with boreal trees⁴⁶. These environments likely expanded during millennial interstadial phases between 59 and 44 ka: GI16, GI14 and GI12⁴⁷, although the analogues for GI12 forest stage (47–44 ka) are both from Central European and Altai-Sayan regions which may suggest an increase in climatic continentality (Supplementary Table S1). In this framework the expression of Heinrich Stadial 5, i.e., 50/49–47.3 ka^{2,48}, is possibly related to a phase of woody biomass decrease with most of the modern analogues recognized in the Sayan montane area (Fig. 5 and Supplementary Table S1). However, the coarser sample resolution and chronological control of this phase in our record prevents its conclusive attribution and correlation.

48% of pollen spectra between 42 and 39 ka (FPD2) finds no analogues among modern Eurasian pollen spectra (see methods; Fig. 4 and Supplementary Table S1), likely because of the very limited number of modern sites with Cichorioideae values as high as 10–30% (only 31 sites out of 5925 modern pollen assemblages). Although this condition prevents any interpretation in terms of modern analogue vegetation structure, some considerations can be made on the basis of fossil pollen data. Fimon PD pollen assemblages share similarities with steppe forb/shrub Plant Functional Type (PFT) described in Ref.⁵³. Several indicator taxa (e.g., Asteraceae such as *Xeranthemum inapertum* type, *Centaurea scabiosa* and *C. nigra*, *Anthemis* type) together with Cichorioideae and Rubiaceae, roughly recall the set of late-flowering herbs blooming today in eastern European meadow steppe (e.g., *Galium verum*, *Centaurea ruthenica*, *Jurinea linearifolia*, *Serratula xeranthemoides*, *Linosyris villosa*)⁵⁴. The scarcity of pollen surface samples in the EMPD2 dataset from the ecozone stretching between Ukraine and Southern Urals may have influenced the MAT results. A step towards drier condition occurred between 40.2 and 38.7 ka: arboreal pollen sank to values < 30%, *Pinus cembra* patches replaced *Pinus sylvestris/mugo* woodlands and xerophytes (*Ephedra* and *Hippophaë*) expanded with *Galium* and *Anthemis* type (Supplementary Fig. S3) indicating sparsely vegetated areas probably exposed to wind erosion and dust deflation. This chronological phase corresponds to HS4 (40.2–38.3 ka)⁵¹.

Between 39 and 27 ka (FPD3–4), the landscape was more diverse than the 60–39 ka interval. It included all types of grasslands, from wet to dry, up to desert-steppe (sensu⁴²) with *Artemisia* and Chenopodiaceae semi-shrubs which are more drought-resistant than grasses, and today replacing them in the transitional region between steppe and semi-desert⁵⁴. The interpretation of an open landscape seems to be in conflict with a relatively large proportion of arboreal pollen, which is over 40% for most of this interval. However, this is largely represented by *Pinus sylvestris/mugo* type, a wind-pollinated conifer which produces extremely high amounts of pollen that can be transported over long distances. Surface pollen spectra in the modern steppe-tundra landscape of Central Asia contain a large proportion of *Pinus* pollen, although *Pinus* is rare or absent from the surrounding landscape⁴¹. These open formations mainly expanded during GS8 to 4 and particularly during HS 3 (30.4 and 29.1 ka)⁵² (Fig. 5). Today such vegetations composed of xerophytes or mesoxerophytes are widespread from the Ural Mountains to the Mongolian Altai in continental, temperate to temperate-cold (subarctic/subalpine) climates where annual precipitation ranges from 400 mm in the north to 200 mm in the south^{55,56}. During interstadials, boreal forests expanded and their tree layer consisted mainly of *Pinus* (*P. sylvestris* and *P. mugo*, although *P. cembra* could also have been present locally) and *Larix* mostly since 31 ka (Fig. 2). Despite the relatively low proportion of *Larix* in pollen spectra, it may have been the dominant tree in most of these forests, as this species is highly underrepresented in pollen spectra due to its low pollen productivity⁵⁷ and limited dispersal. This forest-steppe ecogradient is today characterised by a continental timberline limit (300 mm annual rainfall and boreal continental climate)⁵⁸ with a dry, but not drought period in mid to late summer⁴². The main difference between pollen zones FPD3 and FPD4 is the larger proportion of forests in the latter, pointing to more humid climate conditions⁴². Considering the landscape pattern in the modern Altai-Sayan region and adjacent areas, it is likely that forests were mainly occurring at sites with a more stable moisture supply, i.e., near streams, valley floors and on north-facing slopes⁵⁹.

Fire signal modulation throughout MIS3. The high-resolution sieved charcoal record from Fimon PD highlights enhanced fire activity during/at the boundary of interstadial forest stages, especially those following Heinrich stadials. Most of the local fires are registered during millennia-long interstadials (e.g., GI8 and GI12), while shorter interstadials have a weaker fire signal with the exception of GI4 and GI3 forest stages (see below). Surprisingly, the 60–50 ka interval, also containing long interstadials (e.g., GI16 and 14), diverges from this pattern, registering subdued fire activity (Fig. 5).

Fire activity mirrored the woody biomass rise and decline during millennia-long forest stages (Fig. 5). For example, at the GI8 onset, a sharp rise of woody pollen accumulation rates (from 25 to 120 grains cm⁻² years⁻¹) and charcoal influx (from 0.03 to 0.1 particles cm⁻² years⁻¹) is synchronous at the stratigraphic resolution of the record (i.e., 70 years) between 38,780 and 38,710 years cal BP (Fimon PD modelled mean ages). The adopted resolution matches the duration of stadial-interstadial transitions, estimated to be about 100–200 years⁶⁰, suggesting




Figure 5. Summary plot showing (a) NGRIP $\delta^{18}\text{O}$ record¹⁴⁷ (grey curve), NGRIP temperature reconstruction⁴⁹ (violet curve), and (b) NGRIP dust $[\text{Ca}^{2+}]$ record, plotted on the GICC05 chronology; (c) PAR (Pollen Accumulation Rates; grains $\text{cm}^{-2} \text{year}^{-1}$) of Fimon PD woody taxa and (d) (CHAR) (Charcoal Accumulation rates; charcoal fragments $> 125 \mu\text{m} \text{cm}^{-2} \text{year}^{-1}$) of Fimon PD charcoal plotted on the Fimon chronology (see “Methods” and Supplementary Fig. S4); (e) Ice-Rafted Debris (IRD) record from the Bay of Biscay: sediment core MD04-2845; Sanchez Goñi et al., 2008; age model $< 31.8 \text{ ka cal BP}$ according to Ref.⁵⁰. Grey bands indicate Heinrich Stadials: HS5 (50/49–47.3 ka)^{2,48}; HS4 (40.2–38.3 ka)⁵¹, HS3 (30.4–29.1 ka)⁵².

that local fires occurred concurrently with the warming-induced biomass rise following Heinrich stadials. A different fire signal modulation is recorded during GI4 and GI3, when charcoal peaks are documented at the onset and at the end of interstadial phases, rather than extending to the overall phases of high forest biomass, as observed instead in GI12 and GI8. A different structure in vegetation and in climate transitions between interstadials may account for the observed changes in fire regimes. For example, in the Fimon pollen record boreal woodlands spread during GI4 and GI3, and climate transitions displayed extreme amplitude shifts, up to 12.5 °C in Greenland⁴⁹. The high magnitude signal of GI4 and GI3 fire episodes in the Fimon record may have been influenced by the establishment of a fire-prone palustrine area (ca. 5.7 km^2) for 3.8 ka, i.e., between ca. 31 and 27 ka³². Indeed, since bogs are more vulnerable to burn in early season than other ecosystems, such local conditions may arguably enhanced fuel consumption per fire episode, in turn favouring extensive fires⁶¹. Also, peat deposits are more adapted archives for the registration of individual fire events than lake sediments⁶².

The increase of fire activity during interstadials throughout the Last Glacial Period has already been documented in long marine and terrestrial records^{50,66}. Unfortunately, the low time-resolution of most of the available charcoal records for MIS 3 limits their comparison at a centennial to decadal scale.

Overall, data from Fimon PD record indicate that fire regime in NE-Italy during MIS 3 was mainly driven by changes in biomass, in turn modulated by high-frequency climatic cycles.

Palaeoclimatic and palaeoecological dynamics throughout MIS 3 in NE-Italy. In this section the Fimon PD palaeoecological data are discussed in the light of climatic forcing and integrated with previous studies on small and large mammals, and bird series from archaeological records (Fumane cave, San Bernardino cave and Broion shelter) and open-air contexts in NE-Italy (Settepolesini) during MIS 3. Long interstadials characterized the first part of MIS 3: GI16.1 (58–56.5 ka), GI14 (54.2–49.6 ka) and GI12 (46.8–44.2 ka) (NGRIP

GICC05 chronology), as indicated by isotopic records in the eastern Mediterranean and Europe³. Open cool-temperate mixed woodlands expanded in NE-Italy (Fig. 6) similarly to what documented in other S-European pollen records¹². The presence of open woodlands is also documented by small mammal assemblages uncovered in Mousterian units of the Fumane cave: A11 to A4 dated to 56–44 ka⁶⁷ (Fig. 7); and micromammal associations of Unit II at San Bernardino where *Microtus agrestis* and *Apodemus gr. sylvaticus-flavicollis* are the most abundant species⁶⁸. Forests likely experienced a contraction during HS5, however, this interval is not adequately resolved in the Fimon PD sequence (Fig. 5), but this contraction is known from other S-European and Mediterranean high-resolution pollen records¹². Data from small mammal assemblages in the Mousterian units A7 and A6 older than 45 ka of Fumane cave, support a decrease in woodlands possibly related to HS5⁶⁷. Studies on large mammals in cave and shelters of the Berici Hills (Supplementary SI-2) indicate a prevalence of cervids (e.g., *Cervus elaphus*, *Capreolus capreolus*, *Megaloceros giganteus*) over caprids (*Capra ibex* and *Rupicapra rupicapra*) and bovinds (*Bison priscus* and *Bos primigenius*)^{63,69} and Supplementary SI-3. Estimated relative frequencies trends of change of ungulates between 47 and 41 ka in the Fumane cave show that *Cervus elaphus* reached its maximum between 45 and 44 ka during a phase of cool mixed forests expansion in the Fimon PD record during GI 12 (Fig. 6).

After GI12, a major cooling is centred on GS12 (44.3–43.3 ka)⁴⁷ as documented in NGRIP T reconstruction⁴⁹ (Fig. 5a). Despite this stadial is less prominent in the Atlantic record⁷², its expression is well recognizable over Europe as an extremely cold phase likely occurring over Western Europe, evidenced by deep frost figures in the loess/palaeosol records from northern France at ~ 44–43 ka cal BP⁷³ and Central Europe⁷⁴. In the northern Alps, glaciers advanced between 46.3 and 42.9 ka cal BP and tundra became dominant since 43 ka (Nesseltalgraben, South-eastern Germany^{75,76}). Increasing dust input is recorded in the ELSA dust stack since 44 ka, peaking at ca. 39 ka during HS4⁶⁵ (Fig. 6). The Fimon PD record documents a major decrease in dry boreal forests with grasslands steadily increasing at ca. 44 ka (onset of GS 12, “A” event—Fig. 6). At ca. 42 ka grasslands expanded at the expense of cool-temperate mixed forests (“B” event—Fig. 6) with dry habitats and sparsely vegetated areas peaking at HS4 (“C” event—Fig. 6). These stepped events recalled the main changes in the estimated relative frequencies of ungulates at Fumane cave (Fig. 6b). The abundance of *Cervus elaphus* and *Capreolus capreolus* started to decline at ca. 44 (“A” event in Fig. 6) in favour of *Capra ibex* and *Rupicapra* sp. and further at ca. 42 ka (“B” event in Fig. 6). In the same time interval *Bos* and *Bison* show a quick unimodal trend⁶³. A cold and dry period was indicated by small mammal assemblages in units A3 to A1 (Fumane cave), dated to between 44 and 38.9 ka cal. BP^{27,67}. Alpine ibex, chamois, and marmot were favoured at low altitudes, as well as arctic fox and wolverine. The Po Plain was, in contrast, inhabited by woolly rhinoceros, mammoth, giant deer and bison (Fig. 6 and Supplementary SI-3)^{63,64}.

GI8 is recognizable as a prominent interstadial phase, marked by a major boreal forest expansion in the Fimon PD record (Figs. 2 and 5), which is also observed in most of central and southern-European high-resolution records^{12,18,77} and possibly indicated by small mammal assemblages in units D1c and D1d in the Fumane cave⁶⁷. From GS8 onwards (i.e., since 36.5 ka) NGRIP [Ca²⁺] record indicates increasing dust input frequency in Greenland⁴⁷ (Fig. 5) and major dustiness peaks over Europe has been recorded at 35 and 32 ka (ELSA dust stack⁶⁵), possibly correlated to two major contractions in dry boreal forests at the Fimon site (Fig. 6). Overall, between ca 39–31 ka, in a context of steppe and desert-steppe expansion, cervids declined sharply, while ibex and chamois reached their peak (see Fumane cave)⁶³. In addition to the persistence of the wolverine, all mustelids, wolves and felids (*Panthera leo*, *Panthera pardus*, *Lynx lynx*) show a general increase, accompanied by hyena (*Crocuta crocuta*). Mammoth and giant deer persisted in the Po Plain⁶⁴.

The most recent interval documented in the Fimon PD record, from 31 to 27 ka, is characterized by the occurrence of carnivores (e.g., cave bear), herbivores like moose (*Alces alces*), already documented in previous phases, and also fish and water birds in the cave and shelters of Berici Hills⁷⁸ (Supplementary SI-3). This phase is marked by the GS 5.1 (HS 3) oscillation which produced a major steppe expansion, as documented elsewhere in the southern Alpine foreland (Casaletto Ceredano site⁷⁹), while the short and pronounced interstadials following HS3 led to the renewed expansion of dry boreal forest up to 27.3 ka^{31,32}. The HS3 event is also inferred based on small-mammal remains at the Fumane cave (unit D1e, with no cultural remains) and other sites in Italy, such as Riparo Mochi (first three phases of unit D), Grotta Paglicci (layers 22c-b)⁸⁰ and Broion cave (layers G2-F)⁸¹. A coeval palaeobotanical record from the open-air site of Piovesello (watershed of the Northern Apennine⁸²) also speaks for an arid rocky landscape above the timberline.

Placing the Middle to Upper Palaeolithic transition in an environmental context. The Fimon PD palaeoecological record provides the unique opportunity to discuss the environmental context where last Neanderthals and early Sapiens lived during MIS 3 in NE-Italy. Cultural changes across the Middle to Upper Palaeolithic transition have been acquired from available archaeological sites: three of them were selected due to their extensive radiocarbon chronology and cultural data, namely the Broion shelter, and the Fumane and San Bernardino caves (Figs. 1b, 7).

Neanderthals inhabited NE-Italy in a mosaic of grasslands and forest patches, composed of a mixture of boreal and broad-leaved temperate trees as those occurring today in the Southern Urals, Central-Eastern Europe and central-southern Siberia (Fig. 3). These ecosystems supported cervid populations (e.g., *Cervus elaphus*, *Capreolus capreolus*, *Megaloceros giganteus*), which prevailed over caprids and bovinds as documented in Late Mousterian units at cave and shelters of Berici Hills⁶³. Neanderthals likely dealt with natural fires, e.g., up to 3 local fire events per 1000 years, that could enclose one or more fires each, between 47 and 44 ka (GI12 forest stage; Figs. 5 and 7). In the eastern slope of the Berici Hills close to Lake Fimon, very late Middle Palaeolithic contexts document fireplaces with abundant bones, burnt flints and sparse charcoal particles at Grotta di San Bernardino⁸³ and at the Broion shelter²⁸. Detailed anthracological investigations carried out at the De Nadale cave suggests that, at the




Figure 6. Eco-climatic proxies (from top to bottom): (a) zoological data from the Broion shelter-cave and the San Bernardino Cave for the Berici Hills; from the Fumane cave for the Lessini Mountains⁶³; from Settepolesini for the Po Plain⁶⁴; (b) Estimated diachronic relative frequency of selected (diagnostic) ungulate taxa based on aoristic sums of NISP data computed for 500-year temporal bins in the interval 47.5–40 cal BP⁶³. Dashed lines indicate discontinuous presence of the same taxa before and after the period with quantitative data; (c) Synthetic pollen curves including taxa grouped according to their ecology and climatic preferences, also highlighted by multivariate and correlation analyses (Fig. 3, Supplementary Figs. S5 and S6): Steppe/desert-steppe (*Artemisia*, Chenopodiaceae, *Hippophaë*); Other grasslands including meadow steppe (Cichorioideae, *Anthemis* type, *Aster* type, *Xeranthemum inapertum* type, Ranunculaceae, *Geranium molle* type, *Ephedra fragilis* and *Ephedra distachya* types); Cool-temperate mixed forest (*Alnus glutinosa* type, *Betula*, *Quercus robur/cerris* type, *Tilia*, *Picea*); Dry boreal forest (*Pinus sylvestris/mugo*, *Larix*, *Juniperus*); (d) Central Europe: ELSA Dust Stack-20 terrestrial sediment core, carbonate grain roundness as regional dust proxy⁶³. Light blue rectangles indicate Heinrich Stadials: HS5 (47.3–49/50 ka)^{2,48}; HS4 (40.2–38.3 ka)⁵¹, HS3 (29.1–30.4 ka)⁵².




Figure 7. Comparison between environmental conditions reconstructed from the off-site Fimon PD palaeoecological record covering the 60–27 ka interval and cultural proxies from selected archaeological sites (NE-Italy). The records are plotted on their own time scale. Key to the panels: (a) Fimon PD synthetic pollen record; (b) Sketch of the changes in dominant vegetation formations through time; (c) Fimon PD interpolated charcoal accumulation rates (pieces $\text{cm}^{-2} \text{year}^{-1}$) (black histograms), fire frequency (red line) and statistically significant local fire events plotted as red “+” symbols; (d) Broion shelter, San Bernardino cave (Berici Hills) and Fumane cave (Lessini Mountains) cultural and chronological sequences^{27,69–71}.

onset of MIS 4, Neanderthals preferentially exploited spruce/larch together with pine (*Pinus* sp. *sylvestris*) and birch⁸⁴. A preferential exploitation of larch is also documented in charcoal assemblages and primary combustion features at the Fumane cave (Lessini Mountains; units A5–A6 and A9)^{85,86}, 50 km away from Lake Fimon. The recurrent selection and use of this wood seems to disagree with the sporadic presence/low abundance of larch in the Fimon PD record (Fig. 2). However, we can relate this bias to the underrepresentation of larch in the regional pollen rain (see above). Indeed, this boreal species likely played an important role in the forests at the NE Alpine border, especially at the continental and upper alpine timberlines³¹.

Sapiens at Fumane (Unit A3) and at the Broion shelter (US1g), attested since 44–43 ka BP, experienced an initial phase of increasing woodland opening and grasslands expansion peaking during HS4 (Figs. 6, 7). Between ca. 44–38 ka, climate conditions have been dry enough to promote fires, but that the high degree of landscape openness resulted in very limited or absent local fuel availability for fire to spread (Fig. 7). This vegetation, which finds very few good modern Eurasian analogues, led to the expansion of ecosystems able to sustain large herds of herbivores, particularly Ibex, chamois and large mammals, and their hunters in the Great Adriatic-Po Region (GAPR). The occurrence of woolly rhino (*Coelodonta antiquitatis*), wolverine (*Gulo gulo*) and arctic fox (*Alopex* cfr. *lagopus*) as climatic markers in Uluzzian units further supports this picture. Other hints in this direction come from the Broion cave pollen record from units containing Upper Palaeolithic industries (“He” unit onwards)⁸⁷ younger than ca. 52–42 ka (approximate time range inferred from two old conventional radiocarbon dates from the “I” unit)⁸⁸. Pollen data show a major environmental change marked by the expansion of Cichorioideae-dominated grasslands comparable to findings from the Fimon PD and TdA records (Supplementary Fig. S1). Although it’s well known that pollen deposition in caves is complex and subjected to stochastic elements⁸⁹, a suitable correlation with an “off-site” record corroborates the consistency of the Broion cave record that preserved the

signal of main environmental transformations at a millennial time scale. Similar findings were observed north of the Alps, where early Sapiens lived in a cold steppe-type environment expanding since 43.5 ka (Willendorf II)⁷⁴. The prominent GS 12 stadial phase (44.3–43.3 ka, Fig. 5) may have played a role in triggering these dynamics in S-European ecosystems⁴, although its expression is less remarkable in the Mediterranean⁹⁰. After the Uluzian phase, the Proto-Aurignacian one developed in a context of extremely open and harsh conditions (Fig. 7) roughly corresponding to GS 10-9/HS4, which favoured hunting of alpine ibex and chamois⁹¹. The following Gravettian cultural phase documented at the caves and shelter of the Broion karstic complex between ca. 32.6 and 30.1 ka, is established in widespread steppe/forest-steppe environments, the latter characterized by fire-prone boreal ecosystems^{32,92}, which finds most of their best modern analogues in the Altai-Sayan region (Figs. 4 and 7). Gravettian hunters, who settled the northern edge of the GAPR, were fully adapted to this ecological context. A long-range mobility through this wide alluvial plain, e.g., between sites located over 250 km apart, is attested by petroarchaeological evidence at both Italian and Balkan sides⁹³.

The long-term perspective offered by the present work suggests that Neanderthal and Sapiens locally developed different adaptive strategies to cope with marked changes in their ecosystems and—possibly—with variable demographic pressure over time. Evidence presented here paves the ground for future investigations on the relationship between resource management, technology, and palaeoenvironmental dynamics in NE-Italy.

Methods

Chronology. The Fimon Ponte sulla Debba core (acronym: Fimon PD) chronology was developed for the 21.75–19.39-m interval (Supplementary Fig. S4). We used six ¹⁴C dates made on bulk peat and terrestrial macroremains (wood and charcoal fragments) and one biostratigraphic tie point³⁰. Three radiocarbon dates were obtained from the Fimon PD core and three others from the Fimon Torri di Arcugnano (TdA) core (Table 1). All ages were computed in a single age-depth model after litho-, bio- and chronostratigraphic correlation of the two sequences (see Supplementary Fig. S1 and Ref.³⁷). ¹⁴C dates made on pollen concentrates and bulk samples from the Fimon PD interval between 19.885 and 21.75 m were excluded from our modelling since they mostly yielded age reversals, as previously demonstrated^{30,32}, Supplementary Fig. S1). The tie-point identified at 21.72 m was obtained by tuning the marked rise in *Tilia* and *Picea* pollen curves with the GI 17 onset (i.e., 60.2 ± 0.4 ka³⁸, see Ref.³⁰ for further details). This is in agreement with the well-established evidence of a significant forest recovery at the end of HE 6 in Mediterranean pollen records^{94,95} as well as at Azzano Decimo⁹⁶. Radiocarbon dates were calibrated using the IntCal20 calibration curve⁹⁷ and the age model was calculated using OxCal 4.4 software⁹⁸. We used standard codes and commands in OxCal, including Poisson-Process modelling with variable *k* parameter⁹⁹. Precision was calculated as 95% confidence ranges for the age estimates of each depth (interpolation rate of 0.5 cm⁻¹).

Loss on ignition (LOI). Loss-On-Ignition was performed on 367 volumetric samples from Fimon PD and 178 volumetric samples from Fimon TdA with an automated LECO TGA 601 thermo-gravimetric analyzer (LECO Corporation, USA). Samples were weighted and progressively heated at 105 °C, 550 °C and 980 °C, to estimate water, total organic matter including sulphides (TOM + s), the carbonate fraction (CaCO₃ + S₂^{-x} + SO₄^{-2x}) (including sulphides and sulphates and the residuum¹⁰⁰). The residuum includes siliceous compounds, oxides, and ash deriving from complete charring of the fuel (i.e., inorganic compounds), apart from the carbonatic ash, measured as an aliquot of CaCO₃ (Supplementary Fig. S2).

Magnetic susceptibility. Magnetic susceptibility was measured on 130 points along the Fimon PD and 277 points along the Fimon TdA cores (Fig. 1 and Supplementary Fig. S1) with a Bartington MS2 susceptibility meter equipped with a MS2E sensor. Data were used to characterize stratigraphic units, as a proxy for clastic input and for correlating the two sequences (Fig. S1).

Microbotanical and gradient analysis. 209 samples were analysed for their microbotanical content and prepared using standard methods (including HF and acetolysis) after adding *Lycopodium* tablets for pollen and charcoal concentrations and influx estimates¹⁰¹ at the Lab. of Palynology and Paleocology of CNR-IGAG in Milano. Pollen identification was carried out at the lowest taxonomic level possible at 400×, 630× and 1000× magnifications under a Leica DM-LB light microscope, using atlases^{102–104} and the CNR reference collection. Pollen diagrams were drawn using *Tilia* ver. 2.6.1¹⁰⁵ and Corel Draw X8 for further graphic elaborations. The pollen sum used for % calculations includes trees, shrubs, and all upland herbs. Aquatics and wetland species are excluded. A mean pollen count of 454 ± 248 grains has been reached. Pollen zonation was obtained through constrained incremental sum of squares cluster analysis (Cavalli Sforza's chord distance as dissimilarity coefficient—CONISS¹⁰⁶). Clustering, restricted to taxa whose pollen reached over 2%, is represented by the dendrogram in Fig. 2. A principal component analysis (PCA) was performed on the covariance matrix of log-transformed % selected data (terrestrial pollen taxa > 2%) (Fig. 3). Data standardization and ordination were carried out with the Vegan package¹⁰⁷ in R environment¹⁰⁸. Calculation of correlation coefficients and *p*-values on the same dataset was done in R using the function *rcorr* (package “Hmisc”)¹⁰⁹ and the function *corrplot* (package “corrplot”)¹¹⁰ to plot the correlation matrix (Supplementary Fig. S5–S6). Such analyses allow us to extract the main eco-gradients and to detect correlations between terrestrial taxa in order to refine their ecological grouping, as shown in Figs. 6 and 7.

Modern analogue technique and pollen indicator taxa. The Modern Analogue Technique (MAT) is a widely used statistical approach based on matching individual fossil pollen spectra to a large number of modern pollen assemblages using similarity measures¹¹¹. Given that modern pollen analogues share similar

combinations of taxa originating from comparable plant assemblages, the vegetation parameters of a modern pollen sample can be applied to any fossil sample with similar palynological composition. We used MAT and squared chord distance (SCD) to calculate dissimilarity between pairs of pollen spectra. For eco-regions prediction, only one ‘best analogue’ with the lowest SCD value was selected¹¹². Aquatic and wetlands plants, spores and algae were excluded from the MAT analysis. A selection of the entire EMPD2 dataset (Eurasian Modern Pollen Database vers. 2)³⁹ consisting of 5925 surface samples and 53 modern pollen assemblages from the Southern Urals collected in 2007 were selected and used as a modern calibration set, covering a long climate gradient across Eurasia. We excluded samples located in the Anatolian and Mediterranean biogeographical regions defined according to EEA (<https://www.eea.europa.eu/data-and-maps/data/biogeographical-regions-europe-3>). Most of the herbaceous taxa included in this Eurasian calibration set are harmonized at family or subfamily level (e.g., Asteroideae, Apiaceae, Rubiaceae). This high harmonization level together with an uneven spatial sampling distribution of the modern samples still represents a limitation in the use of MAT and may cause non-analogue conditions. For this reason, climate-sensitive pollen indicators were also used to define past floristic structures and climatic requirements especially for herbaceous-structured vegetations (e.g., *Xeranthemum inapertum* type, *Galium* type).

Analogue computations were performed using the R package ‘rioja’¹¹³. The optimal maximum number of analogues, $k = 4$, was determined by leave-one-out (LOO) cross-validation.

Vegetation data. Modern pollen analogues were assigned to ecoregions based on their geographical coordinates and elevation using the Ecoregions2017 Resolve map⁴³. These analogues were grouped into four time-periods corresponding to pollen zonation: 60–42 ka (pollen zone FPD1), 42–39 ka (FPD2), 39–31 ka (FPD3) and 31–27 ka (FPD 4), and the proportions of ecoregions with the best modern analogues for each pollen zone were calculated (Fig. 4). For the Altai-Sayan region in southern Siberia, where a large proportion of modern analogues was identified, plant species records from plots established at each site where surface pollen was sampled^{15,41,114} were used to interpret the vegetation and landscape of the Fimon PD fossil samples. For surface pollen samples without detailed vegetation descriptions we integrated information using “sample context” and “vegetation description” notes available in the EMPD2 dataset. These data were used to interpret vegetation types and landscape structure throughout MIS 3 at the Fimon site.

Sieved charcoal particles analysis. Three different size ranges (62–125 μm , 125–500 μm and $> 500 \mu\text{m}$) of sieved charcoal particles were separated in 415 sediment samples of approximately 2 cm^3 at contiguous 0.5–2.5 cm intervals using standard sieving methods¹¹⁵. Samples were gently disaggregated in a 50/50 solution of 10% sodium hexametaphosphate (NaPO_3)₆ and 12.5% sodium hypochlorite (NaClO) for 24 h and sieved (62, 125, 500- μm mesh). The sieved fractions were counted on a gridded platform using a stereomicroscope. Macroscopic charcoal particles ($> 125 \mu\text{m}$) are assumed to record high severity fires within a few kilometers from the study site^{115,116}. Finer charcoal particles (62–125 μm size) are more widely dispersed and may represent a reliable proxy of regional fire variability (at least within 150 km) as supported by dispersal models¹¹⁷. We obtained series of charcoal concentrations (particles cm^{-3}) converted into total charcoal accumulation rates (CHAR, particles $\text{cm}^{-2} \text{year}^{-1}$) by multiplying these values by sediment accumulation rates (cm year^{-1}) inferred from the age–depth model (Supplementary Fig. S7). The CHAR record (particles $> 125 \mu\text{m}$) was then decomposed into background ($C_{\text{background}}$) and peak component using the method implemented in CharAnalysis 0.9 software¹¹⁸. Peaks, which are positive deviations from the $C_{\text{background}}$ represent input of charcoal as a result of large, high-severity fires relatively close (1–3 km radii) to coring location, whereas total CHAR correlates best with area burned at distances from 10^0 – 10^1 km, up to an order of magnitude greater than distances defining the optimal spatial scale for peak-inferred fires¹¹⁶. The $C_{\text{background}}$ component was determined using a moving mode robust to outliers with a 500 years window width. A Gaussian mixture model was used to identify threshold values for peak identification (0.95 percentile). The fire frequency (FF) is the total number of fires within a 1000-year window. Fire return interval (FRI) is the time between two adjacent fire events. A Signal to Noise Index (SNI) was used to evaluate the suitability of sediment-charcoal records for reconstructing local fires. The SNI compares the variability in the signal population, $\text{var}(S)$, to the variability in the noise population, $\text{var}(N)$: $\text{SNI} = \text{var}(S)/\text{var}(S) + \text{var}(N)$. A SNI greater than 3 consistently identifies records appropriate for peak detection⁴⁴.

Data availability

All data generated or analysed during this study are included in this published article and its Supplementary Information files.

Received: 3 February 2023; Accepted: 3 July 2023

Published online: 01 August 2023

References

1. Heinrich, H. Origin and consequences of cyclic ice rafting in the Northeast Atlantic Ocean during the past 130,000 years. *Quat. Res.* **29**, 142–152 (1988).
2. Sanchez Goñi, M. F. & Harrison, S. P. Millennial-scale climate variability and vegetation changes during the Last Glacial: Concepts and terminology. *Quat. Sci. Rev.* **29**, 2823–2827 (2010).
3. Moseley, G. E. *et al.* Multi-speleothem record reveals tightly coupled climate between central Europe and greenland during marine isotope stage 3. *Geology* **42**, 1043–1046 (2014).
4. Staubwasser, M. *et al.* Impact of climate change on the transition of Neanderthals to modern humans in Europe. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 9116–9121 (2018).

5. Columbu, A. *et al.* Speleothem record attests to stable environmental conditions during Neanderthal–modern human turnover in southern Italy. *Nat. Ecol. Evol.* (2020) (in press).
6. Allen, J. R. M. *et al.* Rapid environmental changes in southern Europe during the last glacial period. *Nature* **400**, 740–743 (1999).
7. Magri, D. Late Quaternary vegetation history at Lagaccione near Lago di Bolsena (central Italy). *Rev. Palaeobot. Palynol.* **106**, 171–208 (1999).
8. Follieri, M., Magri, D. & Sadori, L. A 250,000-years pollen record from Valle di Castiglione (Roma). *Pollen et spores* **XXX**, 329–356 (1988).
9. Wulf, S. *et al.* The marine isotope stage 1–5 cryptotephra record of Tenaghi Philippon, Greece: Towards a detailed tephrostratigraphic framework for the Eastern Mediterranean region. *Quat. Sci. Rev.* **186**, 236–262 (2018).
10. Margari, V., Gibbard, P. L., Bryant, C. L. & Tzedakis, P. C. Character of vegetational and environmental changes in southern Europe during the last glacial period; evidence from Lesvos Island, Greece. *Quat. Sci. Rev.* **28**, 1317–1339 (2009).
11. Wulf, S., Çağatay, M. N., Appelt, O., Eriş, K. K. & Henry, P. Defining the Upper Nisyros Pumice (57.1 ± 1.5 ka) as new tephra isochrone for linking early MIS-3 palaeoenvironmental records in the Aegean-Black Sea gateway: New evidence from the Sea of Marmara. *Quat. Sci. Rev.* **274**, 107285 (2021).
12. Fletcher, W. J. *et al.* Millennial-scale variability during the last glacial in vegetation records from Europe. *Quat. Sci. Rev.* **29**, 2839–2864 (2010).
13. Badino, F. *et al.* An overview of Alpine and Mediterranean palaeogeography, terrestrial ecosystems and climate history during MIS 3 with focus on the Middle to Upper Palaeolithic transition. *Quat. Int.* **551**, (2020).
14. Tzedakis, P. C., Lawson, I. T., Frogley, M. R., Hewitt, G. M. & Preece, R. C. Buffered tree population changes in a Quaternary refugium: Evolutionary implications. *Science* **1979**(297), 2044–2047 (2002).
15. Kuneš, P. *et al.* Interpretation of the last-glacial vegetation of eastern-central Europe using modern analogues from southern Siberia. *J. Biogeogr.* **35**, 2223–2236 (2008).
16. Magyari, E. K. *et al.* Late Pleniglacial vegetation in eastern-central Europe: Are there modern analogues in Siberia?. *Quat. Sci. Rev.* **95**, 60–79 (2014).
17. Moine, O. *et al.* The impact of Last Glacial climate variability in west-European loess revealed by radiocarbon dating of fossil earthworm granules. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 6209–6214 (2017).
18. Kern, O. A. *et al.* A near-continuous record of climate and ecosystem variability in Central Europe during the past 130 kyr (Marine Isotope Stages 5–1) from Füramoos, southern Germany. *Quat. Sci. Rev.* **284**, (2022).
19. Hublin, J. J. The modern human colonization of western Eurasia: When and where?. *Quat. Sci. Rev.* **118**, 194–210 (2015).
20. Benazzi, S. *et al.* Early dispersal of modern humans in Europe and implications for Neanderthal behaviour. *Nature* **479**, 525–528 (2011).
21. Banks, W. E., d’Errico, F. & Zilhão, J. Human-climate interaction during the Early Upper Paleolithic: Testing the hypothesis of an adaptive shift between the Proto-Aurignacian and the Early Aurignacian. *J. Hum. Evol.* **64**, 39–55 (2013).
22. Higham, T. *et al.* The timing and spatiotemporal patterning of Neanderthal disappearance. *Nature* **512**, 306–309 (2014).
23. Talamo, S. *et al.* The new 14C chronology for the Palaeolithic site of La Ferrassie, France: The disappearance of Neanderthals and the arrival of Homo sapiens in France. *J. Quat. Sci.* **35**, 961–973 (2020).
24. Peresani, M., Bertola, S., Delpiano, D., Benazzi, S. & Romandini, M. The Uluzzian in the north of Italy: Insights around the new evidence at Riparo Broion. *Archaeol. Anthropol. Sci.* **11**, 3503–3536 (2019).
25. Benazzi, S. *et al.* The makers of the Protoaurignacian and implications for Neanderthal extinction. *Science* **1979**(348), 793–796 (2015).
26. Benazzi, S. *et al.* Middle paleolithic and uluzzian human remains from fumane cave, Italy. *J. Hum. Evol.* **70**, 61–68 (2014).
27. Higham, T. *et al.* Problems with radiocarbon dating the Middle to Upper Palaeolithic transition in Italy. *Quat. Sci. Rev.* **28**, 1257–1267 (2009).
28. Romandini, M. *et al.* A late Neanderthal tooth from northeastern Italy. *J. Hum. Evol.* **147**, 102867 (2020).
29. Peresani, M. Inspecting human evolution from a cave. Late Neanderthals and early sapiens at Grotta di Fumane: present state and outlook. *J. Anthropol. Sci. Istituto Italiano di Antropol.* **100**, 71–107 (2022).
30. Pini, R., Ravazzi, C. & Reimer, P. J. The vegetation and climate history of the last glacial cycle in a new pollen record from Lake Fimon (southern Alpine foreland, N-Italy). *Quat. Sci. Rev.* **29**, 3115–3137 (2010).
31. Pini, R. *et al.* Linking North Atlantic and Alpine Last Glacial Maximum climates via a high-resolution pollen-based subarctic forest steppe record. *Quat. Sci. Rev.* **294**, 107759 (2022).
32. Badino, F. *et al.* The fast-acting “pulse” of Heinrich Stadial 3 in a mid-latitude boreal ecosystem. *Sci. Rep.* **10**, 1–14 (2020).
33. Edwards, K., Harris, D. R. & Thomas, K. D. Using space in cultural palynology: The value of the off site pollen record. In *Modeling Ecological Change. Perspectives from Neocology, Palaeoecology and Environmental Archaeology* 61–71 (1991).
34. Pellegrini, C. *et al.* The late Pleistocene Po River lowstand wedge in the Adriatic Sea: Controls on architecture variability and sediment partitioning. *Mar. Pet. Geol.* **96**, 16–50 (2018).
35. Waelbroeck, C. *et al.* Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records. *Quat. Sci. Rev.* **21**, 295–305 (2002).
36. Antonioli, F. *et al.* New evidence of mis 3 relative sea level changes from the messina strait, Calabria (Italy). *Water (Switzerland)* **13**, (2021).
37. Monegato, G., Pini, R., Ravazzi, C., Reimer, P. J. & Wick, L. Correlating Alpine glaciation with Adriatic sea-level changes through lake and alluvial stratigraphy. *J. Quat. Sci.* **26**, 791–804 (2011).
38. Holzkämper, S., Spötl, C. & Mangini, A. High-precision constraints on timing of Alpine warm periods during the middle to late Pleistocene using speleothem growth periods. *Earth Planet Sci. Lett.* **236**, 751–764 (2005).
39. Davis, B. A. S. *et al.* The Eurasian modern pollen database (EMPD). *Version 2*, 1–41 (2020).
40. Simpson, G. L. Analogue methods in palaeolimnology. In *Tracking Environmental Change Using Lake Sediments. Developments in Palaeoenvironmental Research* (Springer, 2012).
41. Pelánková, B. & Chytrý, M. Surface pollen-vegetation relationships in the forest-steppe, taiga and tundra landscapes of the Russian Altai Mountains. *Rev. Palaeobot. Palynol.* **157**, 253–265 (2009).
42. Walter, H., Breckle, S.-W., Walter, H. & Breckle, S.-W. The zonococtones. In *Ecological Systems of the Geobiosphere* 104–107 (Springer, 1986). https://doi.org/10.1007/978-3-662-06812-0_10.
43. Dinerstein, E. *et al.* An ecoregion-based approach to protecting half the terrestrial realm. *Bioscience* **67**, 534–545 (2017).
44. Kelly, R. F., Higuera, P. E., Barrett, C. M. & Sheng Hu, F. A signal-to-noise index to quantify the potential for peak detection in sediment-charcoal records. *Quat. Res.* <https://doi.org/10.1016/j.yqres.2010.07.011> (2011).
45. Chytrý, M. *et al.* A modern analogue of the Pleistocene steppe-tundra ecosystem in southern Siberia. *Boreas* **48**, 36–56 (2019).
46. Chytrý, M. *et al.* Modern analogues from the Southern Urals provide insights into biodiversity change in the early Holocene forests of Central Europe. *J. Biogeogr.* **37**, 767–780 (2010).
47. Rasmussen, S. O. *et al.* A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: Refining and extending the INTIMATE event stratigraphy. *Quat. Sci. Rev.* **106**, 14–28 (2014).
48. Goñi, M. F. S. Regional impacts of climate change and its relevance to human evolution. *Evol. Hum. Sci.* **2**, 1–27 (2020).
49. Kindler, P. *et al.* Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core. *Clim. Past* **10**, 887–902 (2014).

50. Daniau, A. L., Goñi, M. F. S. & Duprat, J. Last glacial fire regime variability in western France inferred from microcharcoal preserved in core MD04-2845, Bay of Biscay. *Quat. Res.* **71**, 385–396 (2009).
51. Cheng, H. *et al.* Onset and termination of Heinrich Stadial 4 and the underlying climate dynamics. *Commun. Earth Environ.* **2**, (2021).
52. Hughen, K. A. & Heaton, T. J. Updated Cariaco Basin C calibration dataset from 0–60 cal kyr BP. *Radiocarbon* **62**, 1001–1043 (2020).
53. Tarasov, P. E. *et al.* A method to determine warm and cool steppe biomes from pollen data; Application to the Mediterranean and Kazakhstan regions. *J. Quat. Sci.* **13**, 335–344 (1998).
54. Walter, H. & Breckle, S.-W. Temperate and polar zonobiomes of Northern Eurasia. In *Ecological Systems of the Geobiosphere* 1–581 (1989).
55. Yurtsev, B. A. The Pleistocene ‘Tundra-steppe’ and the productivity paradox: The landscape approach. *Quat. Sci. Rev.* **20**, 165–174 (2001).
56. Archibold, O. W. *Ecology of World Vegetation. Ecology of World Vegetation* (Springer Netherlands, 1995). <https://doi.org/10.1007/978-94-011-0009-0>.
57. Geng, R. *et al.* Modern pollen assemblages from lake sediments and soil in east siberia and relative pollen productivity estimates for major taxa. *Front. Ecol. Evol.* **10**, 1–17 (2022).
58. Makunina, N. I. Botanical and geographical characteristics of forest steppe of the Altai-Sayan mountain region. *Contemp. Probl. Ecol.* **9**, 342–348 (2016).
59. Hais, M., Chytrý, M. & Horsák, M. Exposure-related forest-steppe: A diverse landscape type determined by topography and climate. *J. Arid Environ.* **135**, 75–84 (2016).
60. Daniau, A. L., Harrison, S. P. & Bartlein, P. J. Fire regimes during the Last Glacial. *Quat. Sci. Rev.* **29**, 2918–2930 (2010).
61. Bourgeau-Chavez, L. L. *et al.* Assessing boreal peat fire severity and vulnerability of peatlands to early season wildland fire. *Front. Genet.* **3**, 1–13 (2020).
62. Remy, C. C. *et al.* Guidelines for the use and interpretation of palaeofire reconstructions based on various archives and proxies. *Quat. Sci. Rev.* **193**, 312–322 (2018).
63. Romandini, M. *et al.* Macromammal and bird assemblages across the late Middle to Upper Palaeolithic transition in Italy: An extended zooarchaeological review. *Quat. Int.* **551**, (2020).
64. Gallini, V. & Sala, B. Settepolesini di Bondeno (Ferrara-Eastern Po Valley): The first example of mammoth steppe in Italy. In *The World of Elephants—International Congress 272–275* (2001).
65. Fuhrmann, F., Seelos, K. & Sirocko, F. Eolian sedimentation in central European Auel dry maar from 60 to 13 ka. *Quat. Res. U.S.* **101**, 4–12 (2021).
66. Magri, D. Lineamenti della vegetazione tardoglaciale in Italia peninsulare e in Sicilia. In *Il Tardiglaciale in Italia—Lavori in corso* 1–8 (2008).
67. López-García, J. M., Dalla Valle, C., Cremaschi, M. & Peresani, M. Reconstruction of the Neanderthal and Modern Human landscape and climate from the Fumane cave sequence (Verona, Italy) using small-mammal assemblages. *Quat. Sci. Rev.* **128**, 1–13 (2015).
68. López-García, J. M., Luzi, E. & Peresani, M. Middle to Late Pleistocene environmental and climatic reconstruction of the human occurrence at Grotta Maggiore di San Bernardino (Vicenza, Italy) through the small-mammal assemblage. *Quat. Sci. Rev.* **168**, 42–54 (2017).
69. Terlato, G. *et al.* Late Neanderthal Subsistence At San Bernardino Cave (Berici Hills—Northeastern Italy) Inferred From Zooarchaeological Data. *Alp. Mediterr. Quat.* **34**, 213–235 (2021).
70. De Stefani, M., Gurioli, F. & Ziggotti, S. Il Paleolitico superiore del Riparo del Broion nei Colli Berici (Vicenza). *Rivista di scienze preistoriche : LV, Supplemento 1* (2005).
71. Douka, K. *et al.* On the chronology of the Uluzzian. *J. Hum. Evol.* **68**, 1–13 (2014).
72. Darfeuille, S. *et al.* Sea surface temperature reconstructions over the last 70 kyr off Portugal: Biomarker data and regional modeling. *Paleoceanography* **31**, 40–65 (2016).
73. Antoine, P. *et al.* Upper Pleistocene loess-palaeosol records from Northern France in the European context: Environmental background and dating of the Middle Palaeolithic. *Quat. Int.* **411**, 4–24 (2016).
74. Nigst, P. R. *et al.* Early modern human settlement of Europe north of the alps occurred 43,500 years ago in a cold steppe-type environment. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 14394–14399 (2014).
75. Stojakowits, P. *et al.* Impact of climatic extremes on Alpine ecosystems during MIS 3. *Quat. Sci. Rev.* **239**, (2020).
76. Mayr, C. *et al.* High-resolution geochemical record of environmental changes during MIS 3 from the northern Alps (Nesseltalgraben, Germany). *Quat. Sci. Rev.* **218**, 122–136 (2019).
77. Badino, F. *et al.* An overview of Alpine and Mediterranean palaeogeography, terrestrial ecosystems and climate history during MIS 3 with focus on the Middle to Upper Palaeolithic transition. *Quat. Int.* <https://doi.org/10.1016/j.quaint.2019.09.024> (2019).
78. Carrera, L. Birds from the late pleistocene: Environmental and climatic scenarios between the Alps and the Great Adriatic Plain (North-Eastern Italy). *Alp. Mediterr. Quat.* **31**, 49–53 (2018).
79. Ravazzi, C. *et al.* Birch-sedge communities, forest withdrawal and flooding at the beginning of Heinrich Stadial 3 at the southern Alpine foreland. *Rev. Palaeobot. Palynol.* **280**, (2020).
80. Berto, C., Boscato, P., Boschin, F., Luzi, E. & Ronchitelli, A. Paleoenvironmental and paleoclimatic context during the Upper Palaeolithic (late Upper Pleistocene) in the Italian Peninsula. The small mammal record from Grotta Paglicci (Rignano Garganico, Foggia, Southern Italy). *Quat. Sci. Rev.* **168**, 30–41 (2017).
81. Berto, C. *Distribuzione ed evoluzione delle associazioni a piccoli mammiferi nella penisola italiana durante il Pleistocene superiore.* (Italian: Università degli Studi di Ferrara, 2013).
82. Peresani, M. *et al.* Human settlement and vegetation-climate relationships in the Greenland Stadial 5 at the Piovesello site (Northern Apennines, Italy). *Quat. Res. U.S.* **90**, 503–528 (2018).
83. Peresani, M. The end of the Middle Palaeolithic in the Italian Alps. An overview on Neanderthal land-use, subsistence and technology. In *Neanderthal Lifeways, Subsistence and Technology—One Hundred Fifty Years of Neanderthal Study* (eds. Conard, N. J. & Richter, J.) 249–259 (Springer, 2011).
84. Vidal-Matutano, P., Livraghi, A. & Peresani, M. New charcoal evidence at the onset of MIS 4: First insights into fuel management and the local landscape at De Nadale cave (northeastern Italy). *Rev. Palaeobot. Palynol.* **298**, 104594 (2022).
85. Basile, D., Castelletti, L. & Peresani, M. Results from the anthracological investigation of the Mousterian layer A9 of Grotta di Fumane, Italy. *Quartar* **61**, 103–111 (2014).
86. Peresani, M. *et al.* Fire-places, frequentations and the environmental setting of the final Mousterian at Grotta di Fumane: A report from the 2006–2008 research. *Quartar* **58**, 131–151 (2011).
87. Cattani, L. & Renault-Miskovsky, J. Etude pollinique du remplissage de la Grotte 1611 du Broion (Vicenza, Italie): Paléoclimatologie du Würmien en Vénétie. *Bull. Assoc. Fr. 1612 Étude Quat.* **XVI**, 197–212 (1984).
88. Leonardi, P. & Broglio, A. Datazione assoluta di un’induztria musteriana della Grotta del Broion. *Rivista di Scienze Preistoriche* **XXI**, 397–405 (1966).
89. Hunt, C. O. & Fiacconi, M. Pollen taphonomy of cave sediments: What does the pollen record in caves tell us about external environments and how do we assess its reliability?. *Quat. Int.* **485**, 68–75 (2018).

90. Vidal-Cordasco, M., Ocio, D., Hickler, T. & Marín-Arroyo, A. B. Ecosystem productivity affected the spatiotemporal disappearance of Neanderthals in Iberia. *Nat. Ecol. Evol.* **6**, (2022).
91. Marín-Arroyo, A. B., Terlato, G., Vidal-Cordasco, M. & Peresani, M. Subsistence of early anatomically modern humans in Europe as evidenced in the Protoaurignacian occupations of Fumane Cave, Italy. *Sci. Rep.* **13**, 1–23 (2023).
92. Pini, R. *et al.* Linking North Atlantic and Alpine Last Glacial Maximum climates via a high-resolution pollen-based subarctic forest steppe record. *Quat. Sci. Rev.* **294**, (2022).
93. Peresani, M. *et al.* Hunter-gatherers across the great Adriatic-Po region during the Last Glacial Maximum: Environmental and cultural dynamics. *Quat. Int.* **581–582**, 128–163 (2021).
94. Tzedakis, P. C. *et al.* Ecological thresholds and patterns of millennial-scale climate variability: The response of vegetation in Greece during the last glacial period. *Geology* **32**, 109–112 (2004).
95. Sánchez Goñi, M. F. *et al.* Contrasting impacts of Dansgaard-Oeschger events over a western European latitudinal transect modulated by orbital parameters. *Quat. Sci. Rev.* **27**, 1136–1151 (2008).
96. Pini, R., Ravazzi, C. & Donegana, M. Pollen stratigraphy, vegetation and climate history of the last 215 ka in the Azzano Decimo core (plain of Friuli, north-eastern Italy). *Quat. Sci. Rev.* **28**, 1268–1290 (2009).
97. Reimer, P. *et al.* The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 kcal BP). *Radiocarbon* <https://doi.org/10.1017/RDC.2020.41> (2020).
98. Bronk Ramsey, C. Bayesian analysis of radiocarbon dates. *Radiocarbon* **51**, 337–360 (2009).
99. Ramsey, C. B. & Lee, S. Recent and planned developments of the program OxCal. *Radiocarbon* **55**, 720–730 (2013).
100. Santisteban, J. I. *et al.* LOI qualitative or quantitative method for organic matter and carbonate mineral content in sediments. (2004).
101. Stockmarr, J. Tablets with spores used in absolute pollen analysis. (1971).
102. Reille, M. Pollen et spores d'Europe et d'Afrique du Nord—Supplément II. (1998).
103. Beug, H. J. *Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete* (Verlag Dr, 2004).
104. Reille, M. *Pollen et Spores d'Europe et d'Afrique du Nord. Laboratoire de Botanique Historique et Palynologie*. (1992).
105. Grimm, E. C. Tilia Version 2.6. 1. Preprint at (2019).
106. Grimm, E. C. CONISS: A Fortran 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares. *Comput. Geosci.* **13**, 13–35 (1987).
107. Oksanen, J. *et al.* Package 'vegan'. *R package version 3.4.0* (2019).
108. R Development Core Team. A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing* <https://www.R-project.org> Preprint at (2019).
109. Harrell, F. & Harrell, M. Package 'Hmisc'. CRAN. 235 Preprint at (2013).
110. Wei, T. *et al.* R package 'corrplot': Visualization of a correlation matrix. *Statistician* **56**, 316–324 (2017).
111. Overpeck, J. T., Webb, T. & Prentice, I. C. Quantitative interpretation of fossil pollen spectra: Dissimilarity coefficients and the method of modern analogs. *Quat. Res.* **23**, 87–108 (1985).
112. Gavin, D. G., Oswald, W. W., Wahle, E. R. & Williams, W. J. A statistical approach to evaluating distance metrics and analog assignments for pollen records. *Quat. Res.* **6**, 356–367 (2003).
113. Juggins, S. rioja: Analysis of Quaternary Science Data. R package version 1.0-5. <https://cran.r-project.org/package=rioja> Preprint at (2022).
114. Chytrý, M. Database of Masaryk University's vegetation research in Siberia. *Biodivers. Ecol.* **4**, 290–290 (2012).
115. Whitlock, C. & Larsen, C. Charcoal as a Fire Proxy. *Track. Environ. Change Using Lake Sedim.* **3**, 75–97 (2002).
116. Higuera, P. E., Whitlock, C. & Gage, J. A. Linking tree-ring and sediment-charcoal records to reconstruct fire occurrence and area burned in subalpine forests of Yellowstone National Park, USA. *Holocene* **21**, 327–341 (2011).
117. Vachula, R. S., Russell, J. M., Huang, Y. & Richter, N. Assessing the spatial fidelity of sedimentary charcoal size fractions as fire history proxies with a high-resolution sediment record and historical data. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **508**, 166–175 (2018).
118. Higuera, P. CharAnalysis 0.9 : Diagnostic and analytical tools for sediment-charcoal analysis. *Montana State University, Bozeman, MT* 1–27 (2009).

Acknowledgements

This work was funded by ERC under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 724046-SUCCESS); website: <https://www.erc-success.eu/>. Thanks are due to Barbora Pelánková for sharing original data on modern pollen assemblages from the Southern Urals and southern Siberia. The authors also thank Paula Reimer and Marzia Breda who provided feedback and critical revision of the manuscript. Thanks are due to all those who contributed to coring and laboratory preparation of the samples. Comments and suggestions by two anonymous reviewers helped improving the manuscript. The authors are indebted to Michela Mariani (Assistant Professor at the University of Nottingham) for her valuable comments and language editing. This paper is a contribution to the CNR-IGAG Research Lines: DTA.AD001.112.001—Studi sui paleoambienti e sul cambiamento climatico nel Quaternario and the DTA. AD001.112.013-SUCCESS-Progetto H2020 ERC-Consolidator Grant Analisi paleoecologiche.

Author contributions

F.B., S.B. (the PI of the ERC n. 724046—SUCCESS), C.R. and R.P. (WP2-Paleoecology collaborators) conceived the idea of the research project. F.B. coordinated the research and wrote the paper with substantial support from R.P., C.R., M.C., M.P., M.R. Pollen, charcoal and LOI analysis were carried out by F.B., R.P. and P.B. Statistical and multivariate analyses were performed by F.B. Crucial ideas towards the overall interpretation of the data were provided by C.R. and R.P. The following contributed to the review of the paper: E.B., L.D. and S.B. All authors contributed critically to the drafts and gave their final approval for publication.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-023-38081-1>.

Correspondence and requests for materials should be addressed to F.B.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

Supplementary information

High-resolution ecosystem changes pacing the millennial climate variability at the Middle to Upper Palaeolithic transition in NE-Italy

Federica Badino^{1,2*}, Roberta Pini², Cesare Ravazzi², Milan Chytrý³, Paolo Bertuletti², Eugenio Bortolini¹, Lydie Dudová⁴, Marco Peresani^{5,2}, Matteo Romandini¹, Stefano Benazzi¹

¹ Department of Cultural Heritage, University of Bologna, 48121 Ravenna, Italy

² CNR - Institute of Environmental Geology and Geoengineering, Lab. of Palynology and Palaeoecology, Research Group on Vegetation, Climate and Human Stratigraphy, 20126 Milano, Italy

³ Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic

⁴ Department of Paleoecology, Institute of Botany, Czech Academy of Sciences, Brno, Czech Republic

⁵ Department of Humanities, Prehistoric and Anthropology Sciences, University of Ferrara, 44100, Ferrara, Italy

* Corresponding author: Federica Badino federica.badino@igag.cnr.it

University of Bologna,

Department of Cultural Heritage

48121 Ravenna, Italy

<http://www.erc-success.eu/>

SI-1

Fimon basin: lithostratigraphical and local environmental setting

Lake Fimon (22 m a.s.l.) is located in a wide basin in the northern sector of the Berici Hills, a karstic plateau 200 km² wide, emerging from the alluvial plain (**Fig. 1**)¹. The Berici Hills belong to the so called Euganeo-Berico-Lessinian wedge², a structural high slightly affected by the Nealpine compression during the Miocene–Quaternary leading to the exhumation of the Berici Hills and bounded to the east by the NW–SE Schio-Vicenza fault³. The plateau is made of Eocene marl and Oligocene limestone, cut by Oligocene mafic volcanic dykes⁴. Younger deposits and residual soils are confined to the karstic sinkholes⁵.

Three cores were drilled through the sedimentary fill of the Fimon Valleys at different locations along a transect from the external sector (open towards the alluvial plain) to the innermost site occupied by Lake Fimon (**Fig. 2c**):

1. Torri di Arcugnano core (Fimon TdA: 48,60 m deep), external sector of the Fimon Valleys (**Fig. 1c and S1**);
 2. Ponte Debba core (Fimon PD: 44.50 m deep) close to the northern lake shore (**Fig. 1c and S1**);
 3. Fimon Lake core (Fimon FL: 16.36 m deep) drilled from the lake surface (**Fig. 1c**).
- Fimon PD and Fimon TdA cores were sampled for pollen analysis, AMS datings, geochemical analysis (LOI), sand petrography, magnetic susceptibility^{6–8} and finely correlated (**Fig. S1**).

The sedimentological structures observed in Fimon PD (21.75 - 19.39 m) and TdA (31 - 26 m) core sections and the high-resolution record of limnic algae and aquatic plants from Fimon PD core (**Fig. S2**) suggest that deposition took place in a shallow lacustrine environment for most of the MIS 3 interval (FL3 lithostratigraphic zone, **Fig. S1-2**). In this context, massive grey clay intervals containing very high siliceous + oxides values (90% of the dry weights) alternate with several organic-rich (silty gyttja) layers marked by flooded littoral zones expansion suggesting a deepening of the water column (FL3b, e, h, g and i; **Fig. S2**). A major variation of the water column height is marked by aquatic algae (p.m.p. *Pediastrum* and *Botryococcus*) and carbonate fraction increase (20.08-20.30 m, FL 3h-FL4; **Fig. S2**). The interval between 19.90-19.39 m is made of peat deposits mainly formed by thin layers of Cyperaceae leaves and characterized by high total organic matter content including sulphides (30-50%; FL5 in **Fig. S2**). The peat interval is attributed to long-lasting

marshy conditions occupying a wide area as recorded by the finding of the same peat layer in the three studied cores^{8,9}. During this phase accumulation rates increased from ca. 0.006 to 0.02 cm/years due to favourable (local) biomass growth under relatively humid conditions¹⁰. A sequence of laminated minerogenic silts rich in limnic algae (FL6; **Fig. S2**) overlaid these terrestrial organic deposits indicate a freshwater input that drowned the mire (see ¹⁰ for further information).

SI-2

Archaeological sites context

Fumane

Grotta di Fumane is located at the foot of the Venetian Prealps in the western Lessini Mountains at an altitude of 350 m and forms a part of a fossil karst system composed of several cavities in dolomitic limestone. This karst complex preserves a well-documented sedimentary sequence 12 m in thickness, divided into four main macro-units based on the lithological features and archaeological remains. From the bottom to top, these four macro-units are distinguished as a sand unit (S), a breccia unit (BR), an anthropogenic unit (A) and a landslide unit (D)¹¹. Macro-unit A, which is the focus of our study, includes several horizontal layers from A13 to A1, ranging from layers dominated by residual dolomite sands (layers A13–A12), angular fine to medium-sized stone layers (levels in the A10 complex, layers A7, A4), stones, slabs and fine material (layer A3) or anthropogenic material (A11, levels in A10 and A9 complexes, A6, A2–A1)¹². Unit A shows evidence of much more intense human occupations, which include (Final) Mousterian (A11, A10 A9, A6–A5, A4), Uluzzian (A3) and Proto-Aurignacian (A2–A1) assemblages^{13–15}. Landslide Unit D mostly consists of large blocks, which collapsed from the cavity's roof because of macrogelivation processes. Evidence of human occupation is present within the lowermost layers D3d, D3b and D3a, representing the latest Aurignacian units ¹⁶ and becoming sporadic in the middle-level D1d, where some Gravettian artefacts have been detected ¹⁷. Research at Fumane is coordinated by University of Ferrara (M.P.) in the framework of a project supported by the Ministry of Culture – “SABAP per le province di Verona, Rovigo e Vicenza”, public institutions (Lessinia Mountain Community - Regional Natural Park, Fumane Municipality, BIMAdige, SERIT) and by private institutions, associations and companies. Research campaigns 2017 and 2019 have received funding from the

European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 724046 – SUCCESS, <http://www.erc-success.eu>).

Broion shelter

Riparo del Broion is situated at 135 m a.s.l. at the base of a steep cliff of Mount Brosimo (327 m a.s.l.) along a terraced slope (Berici Hills). The shelter is 10 m long, 6 m deep and 17 m high and originated from rock collapse along a major ENE-WSW oriented fault that developed from thermoclastic processes and chemical dissolution comparably to other cavities in the area ¹⁸. Two additional Paleolithic cavities were investigated on the western side of the same cliff, Grotta del Buso Doppio del Broion and Grotta del Broion¹⁹.

Archaeological excavations were initially directed by Alberto Broglio (1998 -2008) and by two of us (M.P. and M.R.) in 2015 on a 20sqm area bounded to north and west from the rock walls. Faunal remains and Middle and Upper Paleolithic (Uluzzian, Gravettian and Epigravettian) cultural material was uncovered ²⁰⁻²². The bedrock has not yet been reached. Sediments are mostly small stones and gravel with large prevalence on loams: 16 stratigraphic units planarly bedded have been identified. The lowermost (11, 9, 7 and 4) contain Mousterian artefacts, faunal remains and clearly differentiate in dark-brownish color from the other units.

Research at Riparo Broion is coordinated by the Bologna (M.R.) and Ferrara (M.P.) Universities in the framework of a project supported by the Ministry of Culture – “SABAP per le province di Verona, Rovigo e Vicenza”, public institutions (Longare Municipality), institutions (Leakey Foundation, Spring 2015 Grant; Istituto Italiano di Preistoria e Protostoria). Research campaigns 2017-2019 have received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 724046 – SUCCESS, <http://www.erc-success.eu>).

San Bernardino

Grotta Maggiore di San Bernardino (Mossano, Berici Hills, Vicenza), located at 135 m a.s.l., lies in the foreland of the eastern Italian Alps. The cave opens along the eastern slope of the karst plateau of the Berici Hills on a carbonate sandstone cliff from the Middle Eocene, facing the alluvial plain of the Bacchiglione River and the south-western sector of the Euganean Hills. The cave is 41 m long, 7 m wide and 9 m high and was produced by

thermoclastic processes and chemical dissolution, which widened deep, SE-NW oriented fractures. In the western side of the same slope, a second, shorter cave named Grotta Minore di San Bernardino is situated. The first archaeological excavations were carried out by Prof. P. Leonardi in the 1960s in the area facing the medieval wall, uncovering a Pleistocene sequence with faunal remains and knapped stones^{23,24}. A second cycle of research, between 1986 and 1995, allowed a more detailed reexamination of the different facies of the stratigraphic series between the inner and outer zones of the cave²⁵. At present, the sedimentary sequence is 4.5 m thick and includes eight stratigraphic units with sub-horizontal bedding, which tilt progressively outside the cave. Excavations from 1986 to 1994 confirmed that all the units contained Mousterian industries except unit I, which contained a few Upper Palaeolithic artifacts. Neanderthals used San Bernardino Cave as a place where carcasses processing was finalized, after an initial process at the kill-site, and then prepared for consumption²⁶.

SI-3

Archaeozoological data

Analysis carried out on Late Mousterian levels at Fumane Cave in Lessini Prealps and San Bernardino Cave and Broion Shelter in the Berici Mounts, indicate a prevalence of cervids (e.g., *Cervus elaphus*, *Capreolus capreolus*, *Megaloceros giganteus*) over caprids (*Capra ibex* and *Rupicapra rupicapra*) and bisons (*Bison priscus* and *Bos primigenius*)²⁶⁻³¹. The presence of freshwaters is also evidenced by findings of beaver and moose remains. Afterwards, between ca. 44-39 ka, major ecological changes occurred in faunal assemblages. In stratigraphic reference sequences we notice a consistent decrease of *Cervus elaphus* and *Capreolus capreolus* as opposed to a sharp increase in *Capra ibex* and *Rupicapra rupicapra* (see the Late Mousterian/Uluzzian/ Protoaurignacian transition, at Fumane cave) and the occurrence of cold-adapted bird species currently distributed at high latitude in the Boreal hemisphere^{27,28,30}. The presence of bovids (*Bos/Bison*) is constant, while moose and giant deer are less frequent and well attested in Mousterian and Uluzzian layers of all three sites and in open sites (see Settepolesini³²) during the *Middle Würm*. Wild boar rarer yet, is more abundant at lower elevations (Grotta di San Bernardino, Mousterian Units II + III; Riparo del Broion, Uluzzian layers 1e+1f+1g). The presence of woolly rhinoceros (*Coelodonta antiquitatis*) in the Uluzzian layer A3 at

Fumane Cave and of *Stephanorhinus* sp. at San Bernardino Cave indicate a decreasing temperature trend. This is also supported by the increase in abundance of a variety of carnivores taxa in the Uluzzian unit^{28,33} and, particularly, the presence of wolverine (*Gulo gulo*), ermine (*Mustela erminea*), and arctic fox (*Alopex* cfr. *lagopus*). Mountain hare (*Lepus timidus*) also appeared between lagomorph's²⁸.

Between 39-31 ka (FPD3), the degrowth of cervids becomes a collapse, while the abundance of ibex and chamois reached its *acme* (see Fumane cave). In this chronophase, at Fumane Cave, mountain hare (*Lepus timidus*) is still present^{27,28,30}.

As regards the alluvial plain of the Po River, Prof. Gallini and Sala B. wrote about Settepolesini in the Middle Würm (ca. 35,000 – 33,000 years ago): “*The findings of mammoth and Irish elk, ascribed by the radiocarbon dating's to this period, together with woolly rhinoceros, steppe bison and elk.*”³².

Between 31 to 27 ka, faunal data from caves and shelters around the Berici Hills record the occurrence of *Alces alces*, fishes and water bird remains, pointing to the presence of water bodies. Marmots, ibex and chamois also occurred^{19,29,34–36}. As for birds, boreal species that possibly sought refuge in the Mediterranean areas during colder phases, are documented at Fumane cave³⁰. The subsequent climatic deterioration linked to the LGM initiation at 27.3 ka³⁷ could have been responsible, together with an increasing human pressure, to the later extinction of cave bears in Northeastern Italy^{38,39}.

All these phases are well detailed by the analysis of bird assemblage at Fumane cave. Several identified bird species (rock ptarmigan, black grouse, Boreal owl, bearded vulture, white-backed woodpecker, red and yellow-billed chough, northern nutcracker and white-winged snow finch, currently live in Italy at considerably higher altitudes than Fumane cave. The presence of their fossil remains at 350 m asl suggests the downward shift of the vegetation belts during MIS 3. The presence of remains probably belonging to willow grouse in layer A6 and to parrot crossbill in layers A1+A2 might be an example of two boreal species seeking a *refugium* in Mediterranean Europe^{35,40}.




Figure S1. Litho-, chrono- and biostratigraphic correlation of Fimon PD and Fimon TdA cores, plotted versus their original depth. ¹⁴C ages are plotted according to their stratigraphic position. Radiocarbon ages indicated in red were excluded from our modelling since they mostly yielded age reversals, as previously demonstrated^{8,41}. Magnetic susceptibility, LOI-determined total organic matter including sulphides, CaCO₃ including sulphides and sulphates and residuum, accompanied by the percentage curves of *Pinus sylvestris/mugo* type, *Artemisia* and Cichorioideae were used to correlate the sequences. Also, data shows the reproducibility of the two Lake Fimon records.




Figure S2 - Fimon PD geochemical data and local environments. Normalized Sum1: wet meadows (*Filipendula*, *Thalictrum* and *Lythrum salicaria*); Sum2: mudflates ephemeral herbs (*Cichorioideae* and *Xeranthemum inapertum* type); Sum3: flooded or waterlogged littoral zones (*Anagallis*, *Lythrum portula*, *Sparganium emersum* type, *Ceratophyllum*, *Typha latifolia* type); Sum4: open waters (*Nymphaea*, *Nuphar*, *Myriophyllum verticillatum*, *Myriophyllum spicatum*, *Myriophyllum alterniflorum*, *Spirogyra*, *Spirogyra scrobiculata*, *Zygnemataceae*, *Mougeotia*, *Pediastrum*, *Botryococcus*, *Scenedesmus*, *Gloeotrichia*).




Figure S4. Fimon PD age-depth model calculated with the OxCal 4.4 calibration software⁴² using IntCal20 calibration curve⁴³.




Figure S5. Correlation matrix of Fimon PD woody taxa > 2%.




Figure S6. Correlation matrix of Fimon PD upland herbs > 2%.




Figure S7. From bottom to the top: Macroscopic charcoal accumulation rates (charcoal pieces >125 micron $\text{cm}^{-2}\text{yr}^{-1}$; grey curve); Fire frequency (FF): the total number of fires within a 1000-year window; Fire return interval (FRI, 1000-yr mean) and 95% CI: the time between two adjacent fire events; Peak magnitude ($\text{pieces cm}^{-2} \text{peak}^{-1}$); Local fires are indicated by red crosses (positive passing the charcoal peak screening and $\text{SNI}>3$) while other peaks: grey crosses (positive passing the charcoal peak screening and $\text{SNI}<3$) and grey dots (negative passing the charcoal peak screenin), identified using the method implemented in CharAnalysis 0.9⁴⁴. Histograms show Pollen

Accumulation Rates (PAR) of all pollen taxa and selected pollen taxa (*Pinus sylv./mugo*, *Picea*, *Betula*, *Larix*, *Juniperus*, *Quercus cerris/robur* type + *Ulmus*).




Figure S8. Boxplot of all Signal to Noise (SNI) values⁴⁵. Analysis were carried out using the method implemented in CharAnalysis 0.9 software⁴⁴.

FIMON PD record		METADATA FROM EMPD2						ECOZONES AND BIOMES	
Fossil samples - pollen zones	Fossil samples - modelled mean ages	First analogue - Sample Name	Longitude	Latitude	Elevation	SampleContext	VegDescription	ECOZONE_NAME	BIOME_NAME
FPD4	26752	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD4	26871	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	27001	Kunes_a46	87,80139	50,10583	2066	treeless vegetation	Mesic grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	27047	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	27094	Kunes_a97	85,67556	50,98500	1382	closed forest	Taiga/Pinus sibirica-Picea	Altai montane forest and forest steppe	Temperate Conifer Forests
FPD4	27140	Kunes_a45	87,78972	50,09417	2094	closed forest	Hemiboreal forest/Pinus sibirica	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	27185	Ortu_b49	6,98400	44,68300	2570	0	0	Alps conifer and mixed forests	Temperate Conifer Forests
FPD4	27231	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands

FPD4	27277	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	27323	Binney2017_E2846	104,32000	51,47000	475	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	27369	Binney2017_E2846	104,32000	51,47000	475	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	27415	Binney2017_E2846	104,32000	51,47000	475	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	27464	Binney2017_E2846	104,32000	51,47000	475	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	27515	Ortu_b49	6,98400	44,68300	2570	0	0	Alps conifer and mixed forests	Temperate Conifer Forests
FPD4	27568	Ryabogina_a27	65,54812	56,48371	64	wetland	near-water vegetation (reeds, sedges, duckweed)	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD4	27620	Binney2017_E2863	107,40000	53,20000	480	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	27671	Ortu_b49	6,98400	44,68300	2570	0	0	Alps conifer and mixed forests	Temperate Conifer Forests
FPD4	27724	Binney2017_E2728	104,72000	52,08000	475	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	27776	Binney2017_E3546	129,33000	62,00000	200	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	27828	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	27880	Kunes_a46	87,80139	50,10583	2066	treeless vegetation	Mesic grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	27932	Kunes_a49	87,80000	50,10583	2085	treeless vegetation	Meadow steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands

FPD4	27984	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28036	Binney2017_E3546	129,33000	62,00000	200	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	28088	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28139	Kunes_a49	87,80000	50,10583	2085	treeless vegetation	Meadow steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28191	Kunes_a49	87,80000	50,10583	2085	treeless vegetation	Meadow steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28243	Kunes_a49	87,80000	50,10583	2085	treeless vegetation	Meadow steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28295	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28347	Kunes_a97	85,67556	50,98500	1382	closed forest	Taiga/Pinus sibirica-Picea	Altai montane forest and forest steppe	Temperate Conifer Forests
FPD4	28399	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28451	EPDcoretop_E320	60,75000	56,76667	230	wetland bog	Pinus sylvestris, Betula pubescens	West Siberian taiga	Boreal Forests/Taiga
FPD4	28503	Binney2017_E2807	106,15000	52,52000	449	0	0		
FPD4	28555	EPDcoretop_E1193	11,67833	46,76083	870	wetland bog	Typha, Phragmites, Utricularia	Alps conifer and mixed forests	Temperate Conifer Forests

FPD4	28607	Kunes_a49	87,80000	50,10583	2085	treeless vegetation	Meadow steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28710	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	28863	Binney2017_E2863	107,40000	53,20000	480	0	0	East Siberian taiga	Boreal Forests/Taiga
FPD4	29017	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	29172	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	29326	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	29481	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	29635	Kunes_a20	87,89639	50,17472	1534	treeless vegetation	Dry steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	29789	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	29943	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	30096	Kunes_a151	88,01139	49,94556	2570	treeless vegetation	Dry steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands

FPD4	30250	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	30405	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	30561	Kunes_a49	87,80000	50,10583	2085	treeless vegetation	Meadow steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	30717	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	30875	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD4	31032	Tonkov_o1	23,40618	41,74560	2230	treeless vegetation	Groups of Pinus mugo within patches of herb vegetation	Rodope montane mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	31188	Lapteva_b5	60,84237	56,25898	295	closed forest	coniferous subtaiga. LocalVeg: pine forest	West Siberian taiga	Boreal Forests/Taiga
FPD3	31425	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD3	31504	Novenko_c7	38,59060	53,67060	155	wetland bog	eutrophic peatland on floodplain	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD3	31661	Lapteva_b86	59,03333	54,10000	477	natural grassland	steppe. LocalVeg: herb-bunchgrass steppe	Kazakh forest steppe	Temperate Grasslands, Savannas & Shrublands

FPD3	31819	Karpinska_c20	21,55670	49,99640	227	pasture	Cirsietum rivularis	Carpathian montane forests	Temperate Conifer Forests
FPD3	31977	Lapteva_a60	57,88907	54,89668	451	0	forest-steppe	Urals montane forest and taiga	Boreal Forests/Taiga
FPD3	32132	Novenko_b4	38,60961	53,67365	176	pasture	Northern forest steppe	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD3	32288	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD3	32445	Novenko_b6	37,62833	54,04417	242	closed forest	broad-leaved forest	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD3	32601	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	32758	Kunes_a54	91,86944	51,71028	1221	treeless vegetation	Dry steppe	Sayan Intermontane steppe	Temperate Grasslands, Savannas & Shrublands
FPD3	32916	Ortu_d39	6,55700	44,61300	1067	closed forest	0	Northeast Spain and Southern France Mediterranean forests	Mediterranean Forests, Woodlands & Scrub
FPD3	33073	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	33230	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	33387	Kunes_a49	87,80000	50,10583	2085	treeless vegetation	Meadow steppe	Altai alpine meadow and tundra	Montane Grasslands & Shrublands

FPD3	33623	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	33857	Kunes_a54	91,86944	51,71028	1221	treeless vegetation	Dry steppe	Sayan Intermontane steppe	Temperate Grasslands, Savannas & Shrublands
FPD3	34014	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	34170	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	34327	Kunes_a82	87,89028	50,29556	2216	treeless vegetation	Alpine grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	34484	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	34798	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	34955	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	35111	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands

FPD3	35266	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	35425	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	35583	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	35740	Kunes_a97	85,67556	50,98500	1382	closed forest	Taiga/Pinus sibirica-Picea	Altai montane forest and forest steppe	Temperate Conifer Forests
FPD3	35897	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	36053	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	36209	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	36366	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands

FPD3	36524	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	36760	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	36917	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	37153	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	37231	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	37310	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	37389	Ryabogina_a27	65,54812	56,48371	64	wetland	near-water vegetation (reeds, sedges, duckweed)	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD3	37467	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	37546	EPDcoretop_E1193	11,67833	46,76083	870	wetland bog	Typha, Phragmites, Utricularia	Alps conifer and mixed forests	Temperate Conifer Forests
FPD3	37624	EPDcoretop_E320	60,75000	56,76667	230	wetland bog	Pinus sylvestris, Betula pubescens	West Siberian taiga	Boreal Forests/Taiga

FPD3	37702	Ryabogina_a27	65,54812	56,48371	64	wetland	near-water vegetation (reeds, sedges, duckweed)	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD3	37780	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	37858	Ryabogina_a27	65,54812	56,48371	64	wetland	near-water vegetation (reeds, sedges, duckweed)	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD3	37937	Ryabogina_a27	65,54812	56,48371	64	wetland	near-water vegetation (reeds, sedges, duckweed)	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD3	38015	Ryabogina_a27	65,54812	56,48371	64	wetland	near-water vegetation (reeds, sedges, duckweed)	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD3	38093	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	38249	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	38406	Kunes_a43	87,80472	50,11472	2013	closed forest	Hemiboreal forest/Pinus sibirica-Larix	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	38562	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	38670	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands

FPD3	38707	Ryabogina_a31	66,12862	56,44082	58	open forest	Grass-meadow grass, there are traces of grazing - plantain, goosefoot, wormwood. Woods with birch, an admixture of pine	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD3	38781	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD3	38854	Kunes_a107	91,16028	53,03694	806	treeless vegetation	Meadow steppe	South Siberian forest steppe	Temperate Grasslands, Savannas & Shrublands
FPD3	38926	Ryabogina_a27	65,54812	56,48371	64	wetland	near-water vegetation (reeds, sedges, duckweed)	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD3	38999	Tonkov_o1	23,40618	41,74560	2230	treeless vegetation	Groups of Pinus mugo within patches of herb vegetation	Rodope montane mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	39072	Tonkov_o1	23,40618	41,74560	2230	treeless vegetation	Groups of Pinus mugo within patches of herb vegetation	Rodope montane mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	39145	Lapteva_b86	59,03333	54,10000	477	natural grassland	steppe. LocalVeg: herb-	Kazakh forest steppe	Temperate Grasslands,

							bunchgrass steppe		Savannas & Shrublands
FPD3	39217	Novenko_b6	37,62833	54,04417	242	closed forest	broad-leaved forest	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD3	39254	Dambach_a269	8,17640	49,02110	130	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD3	39326	Tonkov_o1	23,40618	41,74560	2230	treeless vegetation	Groups of Pinus mugo within patches of herb vegetation	Rodope montane mixed forests	Temperate Broadleaf & Mixed Forests
FPD3	39363	Dambach_a279	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD3	39400	Ryabogina_a8	67,03955	57,39380	59	closed forest	Pine forest, cowberry	Western Siberian hemiboreal forests	Temperate Broadleaf & Mixed Forests
FPD2	39437	Dambach_a279	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39473	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39510	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39547	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39583	Dambach_a276	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests

FPD2	39620	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39656	Dambach_a276	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39692	Dambach_a279	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39729	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39766	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39802	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39838	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39874	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39911	Dambach_a279	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39948	Dambach_a368	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	39984	Dambach_a279	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests

FPD2	40021	Novenko_c7	38,59060	53,67060	155	wetland bog	eutrophic peatland on floodplain	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD2	40094	Dambach_a269	8,17640	49,02110	130	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	40166	Dambach_a277	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	40238	Dambach_a277	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	40311	Novenko_b6	37,62833	54,04417	242	closed forest	broad-leaved forest	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD2	40385	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD2	40458	Kunes_a59	87,64611	50,30750	1342	treeless vegetation	Saline grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD2	40531	EPDcoretop_E2005	0,13000	47,28000	28	riverine	cultivated fields, poplar	European Atlantic mixed forests	Temperate Broadleaf & Mixed Forests
FPD2	40604	Dambach_a279	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	40678	Dambach_a277	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	40750	Lapteva_a60	57,88907	54,89668	451	0	forest-steppe	Urals montane forest and taiga	Boreal Forests/Taiga
FPD2	40822	Dambach_a277	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests

FPD2	40895	Novenko_b4	38,60961	53,67365	176	pasture	Northern forest steppe	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD2	40968	Novenko_b4	38,60961	53,67365	176	pasture	Northern forest steppe	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD2	41041	Novenko_c5	38,70250	53,63944	215	fallow	abandoned cropland	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD2	41187	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	41260	Novenko_b4	38,60961	53,67365	176	pasture	Northern forest steppe	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD2	41332	EPDcoretop_E647	21,61667	49,70000	220	ephemeral lake/pond	fields and meadows	Carpathian montane forests	Temperate Conifer Forests
FPD2	41480	Lapteva_b5	60,84237	56,25898	295	closed forest	coniferous subtaiga. LocalVeg: pine forest	West Siberian taiga	Boreal Forests/Taiga
FPD2	41552	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD2	41624	Novenko_c5	38,70250	53,63944	215	fallow	abandoned cropland	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD2	41696	Tonkov_o1	23,40618	41,74560	2230	treeless vegetation	Groups of Pinus mugo within patches of herb vegetation	Rodope montane mixed forests	Temperate Broadleaf & Mixed Forests

FPD2	41769	Dambach_a369	7,93520	49,66130	415	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD2	41842	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD2	41915	EPDcoretop_E647	21,61667	49,70000	220	ephemeral lake/pond	fields and meadows	Carpathian montane forests	Temperate Conifer Forests
FPD2	41988	Lapteva_a60	57,88907	54,89668	451	0	forest-steppe	Urals montane forest and taiga	Boreal Forests/Taiga
FPD2	42061	Tonkov_o1	23,40618	41,74560	2230	treeless vegetation	Groups of Pinus mugo within patches of herb vegetation	Rodope montane mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	42134	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	42338	Dambach_a18	15,38330	48,38330	790	0	0	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	42543	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	42748	Novenko_b4	38,60961	53,67365	176	pasture	Northern forest steppe	East European forest steppe	Temperate Broadleaf & Mixed Forests
FPD1	42951	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	43155	Pidek_a19	23,05778	50,59472	257	open forest	Nemoral mixed forest, with Picea,	Central European mixed forests	Temperate Broadleaf & Mixed Forests

							Abies, Fagus - national park		
FPD1	43359	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	43563	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	43769	Lapteva_b5	60,84237	56,25898	295	closed forest	coniferous subtaiga. LocalVeg: pine forest	West Siberian taiga	Boreal Forests/Taiga
FPD1	43972	Kunes_a87	87,89028	50,30417	2319	treeless vegetation	Tundra/Dryas	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD1	44175	Novenko_c80	32,97980	56,97980	255	fallow	meadow near forest	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	44380	Pidek_a6	23,06694	50,58222	305	pasture	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	44584	Novenko_c78	32,97640	56,45420	246	fallow	meadow near the village	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	44890	Blyakharchuk_a16	89,34345	50,22395	2748	treeless vegetation	High-mountain area Dzhulukul. Herb—sedge-grass steppe near high-mountain lake.	Sayan alpine meadows and tundra	Montane Grasslands & Shrublands

FPD1	45091	Dambach_a279	8,16280	49,02650	140	0	0	Western European broadleaf forests	Temperate Broadleaf & Mixed Forests
FPD1	45398	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	45804	Pidek_a19	23,05778	50,59472	257	open forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	46315	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	46723	Pidek_a19	23,05778	50,59472	257	open forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	47027	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	47232	Kolaczek_i30	22,44000	54,33000	187	closed forest	LV: Picea abies, Sphagnum sp., Vaccinium myrtillus	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	47437	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	47745	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	47847	Lapteva_b86	59,03333	54,10000	477	natural grassland	steppe. LocalVeg: herb-bunchgrass steppe	Kazakh forest steppe	Temperate Grasslands, Savannas & Shrublands
FPD1	48051	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests

FPD1	48457	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	48865	Lapteva_a53	58,84255	54,52010	1582	forest undefined	boreal forests	Urals montane forest and taiga	Boreal Forests/Taiga
FPD1	49171	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	49480	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	49681	Kunes_a137	92,21306	52,24278	2013	treeless vegetation	Alpine grassland	Sayan montane conifer forests	Temperate Conifer Forests
FPD1	49886	Lapteva_b86	59,03333	54,10000	477	natural grassland	steppe. LocalVeg: herb-bunchgrass steppe	Kazakh forest steppe	Temperate Grasslands, Savannas & Shrublands
FPD1	50089	Kunes_a82	87,89028	50,29556	2216	treeless vegetation	Alpine grassland	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD1	50496	Lapteva_a2	61,24602	54,67487	244	0	forest-steppe	Kazakh forest steppe	Temperate Grasslands, Savannas & Shrublands
FPD1	50698	EPDcoretop_E647	21,61667	49,70000	220	ephemeral lake/pond	fields and meadows	Carpathian montane forests	Temperate Conifer Forests
FPD1	50902	Ortu_b45	6,97800	44,68800	2633	0	0	Alps conifer and mixed forests	Temperate Conifer Forests
FPD1	51413	Pidek_a19	23,05778	50,59472	257	open forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	51928	Lapteva_a2	61,24602	54,67487	244	0	forest-steppe	Kazakh forest steppe	Temperate Grasslands,

									Savannas & Shrublands
FPD1	52437	Pidek_a19	23,05778	50,59472	257	open forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	52945	Kunes_a90	87,88861	50,30583	2499	treeless vegetation	Open scree	Altai alpine meadow and tundra	Montane Grasslands & Shrublands
FPD1	53968	Pidek_a6	23,06694	50,58222	305	pasture	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	54988	Pidek_a5	23,06306	50,58694	279	closed forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	55498	Pidek_a19	23,05778	50,59472	257	open forest	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	56010	Kolaczek_i30	22,44000	54,33000	187	closed forest	LV: Picea abies, Sphagnum sp., Vaccinium myrtillus	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	56723	EPDcoretop_E2327	38,59000	53,67000	161	mire	agricultural land	East European forest steppe	Temperate Broadleaf & Mixed Forests

FPD1	57130	Novenko_c45	40,94190	55,13850	93	fallow	abandoned cropland with small pine	Sarmatic mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	57434	Kolaczek_i30	22,44000	54,33000	187	closed forest	LV: Picea abies, Sphagnum sp., Vaccinium myrtillus	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	58147	EPDcoretop_E375	17,39028	52,55694	109	lake	cultivated fields, deforested area	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	58559	Pidek_a6	23,06694	50,58222	305	pasture	Nemoral mixed forest, with Picea, Abies, Fagus - national park	Central European mixed forests	Temperate Broadleaf & Mixed Forests
FPD1	59068	Lapteva_b86	59,03333	54,10000	477	natural grassland	steppe. LocalVeg: herb-bunchgrass steppe	Kazakh forest steppe	Temperate Grasslands, Savannas & Shrublands
FPD1	59580	Lapteva_b106	58,76667	52,91667	329	cave	steppe	Kazakh forest steppe	Temperate Grasslands, Savannas & Shrublands
FPD1	60090	Lapteva_b106	58,76667	52,91667	329	cave	steppe	Kazakh forest steppe	Temperate Grasslands, Savannas & Shrublands

Table S1 - Coordinates and metadata for the best (first) modern analogues of Fimon PD fossil pollen spectra. Low similarity/non analogue situations are highlighted in bold type.




Figure S9 – Comparison between Fimon PD pollen data: sum of grasslands (Cichorioideae, *Anthemis* type, *Aster* type, *Xeranthemum inapertum* type, *Ranunculaceae*, *Geranium molle* type, *Ephedra fragilis* type, *Ephedra distachya* type); steppe (*Artemisia*, *Chenopodiaceae*, *Hippophae*) and squared-chord distance curves for the calibration set, distances smaller than the 5th percentile of all distances between the calibration set samples are "good analogues", while distances larger than the 10th percentile are "non-analogue" assemblages⁴⁶.

References

1. Fontana, A., Mozzi, P. & Bondesan, A. Alluvial megafans in the Venetian-Friulian Plain (north-eastern Italy): Evidence of sedimentary and erosive phases during Late Pleistocene and Holocene. *Quaternary International* **189**, 71–90 (2008).
2. Massironi, M., Zampieri, D. & Caporali, A. Miocene to present major fault linkages through the Adriatic indenter and the Austroalpine-Penninic collisional wedge (Alps of NE Italy). *Geological Society Special Publication* **262**, 245–258 (2006).
3. Pola, M., Ricciato, A., Fantoni, R., Fabbri, P. & Zampieri, D. Architecture of the western margin of the North Adriatic foreland: the Schio-Vicenza fault system. *Italian Journal of Geoscience* **133**, 223–234 (2014).

4. Macera, P. *et al.* Geodynamic implications of deep mantle upwelling in the source of Tertiary volcanics from the Veneto region (South-Eastern Alps). *Journal of Geodynamics* **36**, 563–590 (2003).
5. Cavinato, A. L'eolico nei M. Berici e Lessini. *Accademia nazionale dei Lincei* (1963).
6. Pini, R., Ravazzi, C. & Reimer, P. J. The vegetation and climate history of the last glacial cycle in a new pollen record from Lake Fimon (southern Alpine foreland, N-Italy). *Quaternary Science Reviews* **29**, 3115–3137 (2010).
7. Monegato, G., Pini, R., Ravazzi, C., Reimer, P. J. & Wick, L. Correlating Alpine glaciation with Adriatic sea-level changes through lake and alluvial stratigraphy. *J Quat Sci* **26**, 791–804 (2011).
8. Badino, F. *et al.* The fast-acting “pulse” of Heinrich Stadial 3 in a mid-latitude boreal ecosystem. *Sci Rep* **10**, 1–14 (2020).
9. Monegato, G., Pini, R., Ravazzi, C., Reimer, P. J. & Wick, L. Correlating Alpine glaciation with Adriatic sea-level changes through lake and alluvial stratigraphy. *Journal of Quaternary Science* **26**, 791–804 (2011).
10. Pini, R. *et al.* Linking North Atlantic and Alpine Last Glacial Maximum climates via a high-resolution pollen-based subarctic forest steppe record. *Quat Sci Rev* **294**, (2022).
11. Abu Zeid, N., Bignardi, S., Russo, P. & Peresani, M. Deep in a Paleolithic archive: Integrated geophysical investigations and laser-scanner reconstruction at Fumane Cave, Italy. *Journal of Archaeological Science: Reports* **27**, 101976 (2019).
12. Peresani, M. *et al.* Age of the final Middle Palaeolithic and Uluzzian levels at Fumane Cave, Northern Italy, using 14C, ESR, 234U/230Th and thermoluminescence methods. *Journal of Archaeological Science* **35**, 2986–2996 (2008).
13. Peresani, M. Fifty thousand years of flint knapping and tool shaping across the Mousterian and Uluzzian sequence of Fumane cave. *Quaternary International* **247**, 125–150 (2012).
14. Peresani, M., Cristiani, E. & Romandini, M. The Uluzzian technology of Grotta di Fumane and its implication for reconstructing cultural dynamics in the Middle-Upper Palaeolithic transition of Western Eurasia. *J Hum Evol* **91**, 36–56 (2016).
15. Falcucci, A., Conard, N. J. & Peresani, M. *A critical assessment of the Protoaurignacian lithic technology at Fumane Cave and its implications for the definition of the earliest Aurignacian.* *PLoS ONE* vol. 12 (2017).
16. Falcucci, A., Conard, N. J. & Peresani, M. Breaking through the aquitaine frame: A re-evaluation on the significance of regional variants during the Aurignacian as seen from a key record in southern Europe. *Journal of Anthropological Sciences* **98**, 99–140 (2020).

17. Falcucci, A. & Peresani, M. A pre-Heinrich Event 3 assemblage at Fumane Cave and its contribution for understanding the beginning of the Gravettian in Italy: Ein vor das Heinrich 3. *Qartär* **66**, 135–154 (2019).
18. Sauro, U. The Monti Berici: A Peculiar Type of Karst in the Southern Alps. *Acta Carsologica* **31**, 99–114 (2016).
19. Romandini, M., Bertola, S. & Nannini, N. Nuovi dati sul Paleolitico dei Colli Berici : risultati preliminari dello studio archeozoologico e delle materie prime litiche della Grotta del Buso Doppio del Broion (Lumignano, Longare, Vicenza). in *Preistoria e protostoria del Veneto. - (Studi di preistoria e protostoria ; 2)* 53–59 (2015).
20. De Stefani, M., Gurioli, F. & Ziggioni, S. Il Paleolitico superiore del Riparo del Broion nei Colli Berici (Vicenza). *Rivista di scienze preistoriche : LV, Supplemento 1* (2005).
21. Peresani, M., Bertola, S., Delpiano, D., Benazzi, S. & Romandini, M. The Uluzzian in the north of Italy: insights around the new evidence at Riparo Broion. *Archaeol Anthropol Sci* **11**, 3503–3536 (2019).
22. Romandini, M. *et al.* A late Neanderthal tooth from northeastern Italy. *J Hum Evol* **147**, 102867 (2020).
23. Leonardi, P. & Broglio, A. Le Paléolithique de la Vénétie. *Annali Università di Ferrara* **15**, 1–118 (1962).
24. Leonardi, P. Risultati della prima campagna di scavo della stazione musteriana di S. Bernardino nei Colli Berici (VI). *Atti Istituto Veneto SS.LL.AA.* **117**, 387–402 (1959).
25. Peresani, M. Sistemi tecnici di produzione litica nel Musteriano d'Italia. Studio tecnologico degli insiemi litici delle unità VI e II della Grotta di San Bernardino (Colli Berici, Veneto). *Rivista Scienze Preistoriche* **47**, 79–167 (1996).
26. Terlato, G. *et al.* Late Neanderthal Subsistence At San Bernardino Cave (Berici Hills - Northeastern Italy) Inferred From Zooarchaeological Data. *Alpine and Mediterranean Quaternary* **34**, 213–235 (2021).
27. Fiore, I., Gala, M. & Tagliacozzo, A. Ecology and subsistence strategies in the Eastern Italian Alps during the Middle Palaeolithic. *International Journal of Osteoarchaeology* **14**, 273–286 (2004).
28. Romandini, M. *et al.* Macromammal and bird assemblages across the late Middle to Upper Palaeolithic transition in Italy: an extended zooarchaeological review. *Quaternary International* **551**, (2020).
29. Cassoli, P. F. & Tagliacozzo, A. I resti ossei di macromammiferi, uccelli e pesci della Grotta maggiore di San Bernardino sui Colli Berici (VI): considerazioni paleoeconomiche, paleoecologiche e cronologiche. *Bull. di Paletnol. Ital* **85**, 1–71 (1994).
30. Cassoli, P. F. & Tagliacozzo, A. Considerazioni paleontologiche, paleoecologiche e archeozoologiche sui macromammiferi e gli uccelli dei livelli del Pleistocene superiore del Riparo di Fumane (VR) scavi 1988-91. *Boll. Museo Civico di Storia Naturale di Verona* **23**, 85–117 (1994).

31. Terlato, G. *et al.* Chronological and Isotopic data support a revision for the timing of cave bear extinction in Mediterranean Europe. *Historical Biology* **31**, 474–484 (2019).
32. Gallini, V. & Sala, B. Settepolesini di Bondeno (Ferrara - Eastern Po Valley): the first example of mammoth steppe in Italy. *The World of Elephants - International Congress 272–275* (2001).
33. Romandini, M. *et al.* Bears and humans, a Neanderthal tale. Reconstructing uncommon behaviors from zooarchaeological evidence in southern Europe. *Journal of Archaeological Science* **90**, 71–91 (2018).
34. Gurioli, F., Cappato, N., De Stefani, M. & Tagliacozzo, A. Considerazioni Paleontologiche, Paleoecologiche e Archeozoologiche dei livelli del Paleolitico superiore del Riparo del Broion (Colli Berici, Vicenza). *Atti del V Convegno Nazionale di Archeozoologia* 47–56 (2006).
35. Carrera, L., Pavia, M., Peresani, M. & Romandini, M. Late Pleistocene fossil birds from Buso Doppio del Broion Cave (North-Eastern Italy): implications for palaeoecology, palaeoenvironment and palaeoclimate. *Bollettino della Società Paleontologica Italiana* **57**, 145–174 (2018).
36. Nannini, N. & Romandini, M. Gravettian and Epigravettian hunters in the Berici Hills (Vicenza): two case of cave bear (*Ursus spelaeus*) exploitation. Cacciatori gravettiani ed epigravettiani nei Colli Berici (Vicenza): due casi di sfruttamento dell'orso delle caverne (*Ursus spelaeus*). *Atti del 7° Convegno Nazionale di Archeozoologia a cura di U. Thun Hohenstein* **11**, (2015).
37. Pini, R. *et al.* Linking North Atlantic and Alpine Last Glacial Maximum climates via a high-resolution pollen-based subarctic forest steppe record. *Quat Sci Rev* **294**, (2022).
38. Gretzinger, J. *et al.* Large-scale mitogenomic analysis of the phylogeography of the Late Pleistocene cave bear. *Sci Rep* **9**, (2019).
39. Terlato, G. *et al.* Extinction and paleoecology of the late pleistocene cave bear from northeastern Italy: Radiocarbon and stable isotope evidence. *Alpine and Mediterranean Quaternary* **31**, 99–103 (2018).
40. Carrera, L. Birds from the late pleistocene: Environmental and climatic scenarios between the Alps and the Great Adriatic Plain (North-Eastern Italy). *Alpine and Mediterranean Quaternary* **31**, 49–53 (2018).
41. Pini, R., Ravazzi, C. & Reimer, P. J. The vegetation and climate history of the last glacial cycle in a new pollen record from Lake Fimon (southern Alpine foreland, N-Italy). *Quat Sci Rev* **29**, 3115–3137 (2010).
42. Bronk Ramsey, C. Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* **51**, 337–360 (2009).
43. Reimer, P. *et al.* The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0-55 kcal BP). *Radiocarbon* 1–33 (2020) doi:doi:10.1017/RDC.2020.41.

44. Higuera, P. CharAnalysis 0.9 : Diagnostic and analytical tools for sediment-charcoal analysis. *Montana State University, Bozeman, MT* 1–27 Preprint at (2009).
45. Kelly, R. F., Higuera, P. E., Barrett, C. M. & Sheng Hu, F. A signal-to-noise index to quantify the potential for peak detection in sediment-charcoal records. (2011) doi:10.1016/j.yqres.2010.07.011.
46. Simpson, G. L. Analogue Methods in Palaeolimnology. in *Tracking Environmental Change Using Lake Sediments. Developments in Paleoenvironmental Research* (Springer, 2012).