

## COLEOPTERA (brouci)

čeled' (podčeled')	taxon	imágo (kód krabice)	larva (číslo epruvety)
<b>Sphaeriusidae</b>	<i>Sphaerius acaroides</i> Waltl, 1838	<b>B1</b>	
<b>Gyrinidae</b>	<i>Gyrinus caspius</i> Ménériés, 1832	<b>B1</b>	
	<i>Gyrinus distinctus</i> Aubé, 1838	<b>B1</b>	
	<i>Gyrinus marinus</i> Gyllenhal, 1808	<b>B1</b>	
	<i>Gyrinus natator</i> (Linnaeus, 1758)	<b>B1</b>	
	<i>Gyrinus paykulli</i> Ochs, 1927	<b>B1</b>	
	<i>Gyrinus substriatus</i> Stephens, 1828	<b>B1</b>	
	<i>Gyrinus</i> sp.		<b>18</b>
	<i>Orectochilus villosus</i> (O. F. Muller, 1776)	<b>B1</b>	<b>19</b>
<b>Haliplidae</b>	<i>Haliplus confinis</i> Stephens, 1828	<b>B1</b>	
	<i>Haliplus flavicollis</i> Sturm, 1834	<b>B1</b>	
	<i>Haliplus fluviatilis</i> Aubé, 1836	<b>B1</b>	
	<i>Haliplus fulvus</i> (Fabricius, 1801)	<b>B1</b>	
	<i>Haliplus heydeni</i> Wehncke, 1875	<b>B1</b>	
	<i>Haliplus immaculatus</i> Gerhardt, 1877	<b>B1</b>	
	<i>Haliplus laminatus</i> (Schaller, 1783)	<b>B1</b>	
	<i>Haliplus lineatocollis</i> (Marsham, 1802)	<b>B1</b>	
	<i>Haliplus obliquus</i> (Fabricius, 1787)	<b>B1</b>	
	<i>Haliplus ruficollis</i> (De Geer, 1774)	<b>B1</b>	
	<i>Haliplus wehncke</i> Gerhardt, 1877	<b>B1</b>	
	<i>Haliplus</i> sp.		<b>20</b>
	<i>Peltodytes caesus</i> (Duftschmid, 1805)	<b>B1</b>	<b>71</b>
	<b>Noteridae</b>	<i>Noterus clavicornis</i> (De Geer, 1774)	<b>B1</b>
<i>Noterus crassicornis</i> (O. F. Müller, 1776)		<b>B1</b>	
<i>Noterus</i> sp.			<b>21</b>
<b>Dytiscidae</b>			
(Hydroporinae)	<i>Bidessus grossepunctatus</i> Vorbringer, 1907	<b>B2</b>	
	<i>Bidessus nasutus</i> Sharp, 1887	<b>B2</b>	
	<i>Bidessus unistriatus</i> (Schrank, 1781)	<b>B2</b>	
	<i>Deronectes latus</i> (Stephens, 1829)	<b>B2</b>	
	<i>Deronectes platynotus</i> (Germar, 1834)	<b>B2</b>	
	<i>Deronectes</i> sp.		<b>22</b>
	<i>Graptodytes granularis</i> (Linnaeus, 1767)	<b>B2</b>	
	<i>Graptodytes pictus</i> (Fabricius, 1787)	<b>B2</b>	
	<i>Hydroglyphus geminus</i> (Fabricius, 1792)	<b>B2</b>	
	<i>Hydroporus angustatus</i> Sturm, 1835	<b>B3</b>	
	<i>Hydroporus erythrocephalus</i> (Linnaeus, 1758)	<b>B3</b>	
	<i>Hydroporus ferrugineus</i> Stephens, 1828	<b>B3</b>	
	<i>Hydroporus gyllenhalii</i> Schiodte, 1841	<b>B3</b>	
	<i>Hydroporus incognitus</i> Sharp, 1869	<b>B3</b>	
	<i>Hydroporus longicornis</i> Sharp, 1871	<b>B3</b>	
	<i>Hydroporus marginatus</i> (Duftschmid, 1805)	<b>B3</b>	
	<i>Hydroporus melanarius</i> Sturm, 1835	<b>B3</b>	
	<i>Hydroporus memnonius</i> Nicolai, 1822	<b>B3</b>	
	<i>Hydroporus nigrita</i> (Fabricius, 1792)	<b>B3</b>	
	<i>Hydroporus obscurus</i> Sturm, 1835	<b>B3</b>	
	<i>Hydroporus palustris</i> (Linnaeus, 1761)	<b>B3</b>	
	<i>Hydroporus planus</i> (Fabricius, 1781)	<b>B3</b>	
	<i>Hydroporus pubescens</i> (Gyllenhal, 1808)	<b>B3</b>	
	<i>Hydroporus sabaudus</i> Fauvel, 1865	<b>B3</b>	
	<i>Hydroporus scalesianus</i> Stephens, 1828	<b>B3</b>	

	<i>Hydroporus striola</i> (Gyllenhal, 1826)	<b>B3</b>	
	<i>Hydroporus tristis</i> (Paykull, 1798)	<b>B3</b>	
	<i>Hydroporus umbrosus</i> (Gyllenhal, 1808)	<b>B3</b>	
	<i>Hydroporus</i> sp.		<b>23</b>
	<i>Hydrovatus cuspidatus</i> (Kunze, 1818)	<b>B2</b>	
	<i>Hygrotus confluens</i> (Fabricius, 1787)	<b>B2</b>	
	<i>Hygrotus decoratus</i> (Gyllenhal, 1810)	<b>B2</b>	
	<i>Hygrotus enneagrammus</i> (Ahrens, 1833)	<b>B2</b>	
	<i>Hygrotus impressopunctatus</i> (Schaller, 1783)	<b>B2</b>	
	<i>Hygrotus inaequalis</i> (Fabricius, 1776)	<b>B2</b>	
	<i>Hygrotus nigrolineatus</i> (Steven, 1808)	<b>B2</b>	
	<i>Hygrotus versicolor</i> (Schaller, 1783)	<b>B2</b>	
	<i>Hygrotus</i> sp.		<b>60</b>
	<i>Hyphydrus ovatus</i> (Linnaeus, 1761)	<b>B2</b>	<b>24</b>
	<i>Nebrioporus canaliculatus</i> (Lacordaire, 1835)	<b>B2</b>	
	<i>Nebrioporus depressus</i> (Fabricius, 1775)	<b>B2</b>	
	<i>Nebrioporus</i> sp.		<b>61</b>
	<i>Oreodytes sanmarkii</i> (C. R. Sahlberg, 1826)	<b>B2</b>	
	<i>Oreodytes septentrionalis</i> (Gyllenhal, 1827)	<b>B2</b>	
	<i>Porhydrus lineatus</i> (Fabricius, 1775)	<b>B2</b>	
	<i>Scarodytes halensis</i> (Fabricius, 1787)	<b>B2</b>	
	<i>Stictotarsus duodecimpustulatus</i> (Fabricius, 1792)	<b>B2</b>	<b>62</b>
	<i>Suphrodytes dorsalis</i> (Fabricius, 1787)	<b>B2</b>	
	Hydroporinae larvae indet.		<b>63</b>
(Laccophilinae)	<i>Laccophilus hyalinus</i> (De Geer, 1774)	<b>B4</b>	
	<i>Laccophilus minutus</i> (Linnaeus, 1758)	<b>B4</b>	
	<i>Laccophilus poecilus</i> Klug, 1834	<b>B4</b>	
	<i>Laccophilus</i> sp.		<b>25</b>
(Copelatinae)	<i>Copelatus haemorrhoidalis</i> (Fabricius, 1787)	<b>B4</b>	
(Colymbetinae)	<i>Agabus affinis</i> (Paykull, 1798)	<b>B5</b>	
	<i>Agabus biguttatus</i> (Olivier, 1795)	<b>B5</b>	
	<i>Agabus bipustulatus</i> (Linnaeus, 1767)	<b>B5</b>	
	<i>Agabus congener</i> (Thunberg, 1794)	<b>B5</b>	
	<i>Agabus guttatus</i> (Paykull, 1798)	<b>B5</b>	
	<i>Agabus labiatus</i> (Brahm, 1790)	<b>B5</b>	
	<i>Agabus melanarius</i> Aubé, 1836	<b>B5</b>	
	<i>Agabus nebulosus</i> (Forster, 1771)	<b>B5</b>	
	<i>Agabus paludosus</i> (Fabricius, 1801)	<b>B5</b>	
	<i>Agabus sturmii</i> (Gyllenhal, 1808)	<b>B5</b>	
	<i>Agabus uliginosus</i> (Linnaeus, 1761)	<b>B5</b>	
	<i>Agabus undulatus</i> (Schränk, 1776)	<b>B5</b>	<b>64</b>
	<i>Agabus unguicularis</i> C.G. Thomson, 1867	<b>B5</b>	
	<i>Agabus</i> sp.		<b>26</b>
	<i>Colymbetes fuscus</i> (Linnaeus, 1758)	<b>B4</b>	<b>27</b>
	<i>Colymbetes striatus</i> (Linnaeus, 1758)	<b>B4</b>	
	<i>Ilybius aenescens</i> C. G. Thomson, 1870	<b>B5</b>	
	<i>Ilybius ater</i> (De Geer, 1774)	<b>B5</b>	
	<i>Ilybius crassus</i> C.G. Thomson, 1854	<b>B5</b>	
	<i>Ilybius fenestratus</i> (Fabricius, 1781)	<b>B5</b>	
	<i>Ilybius fuliginosus</i> (Fabricius, 1792)	<b>B5</b>	
	<i>Ilybius guttiger</i> (Gyllenhal, 1808)	<b>B5</b>	
	<i>Ilybius chalconatus</i> (Panzer, 1796)	<b>B5</b>	
	<i>Ilybius quadriguttatus</i> (Boisduval & Lacordaire, 1835)	<b>B5</b>	
	<i>Ilybius</i> sp.		<b>28</b>
	<i>Platambus maculatus</i> (Linnaeus, 1758)	<b>B4</b>	<b>29</b>
	<i>Rhantus bistriatus</i> (Bergstrasser, 1778)	<b>B4</b>	

	<i>Rhantus consputus</i> (Sturm, 1834)	<b>B4</b>	
	<i>Rhantus exsoletus</i> (Forster, 1771)	<b>B4</b>	
	<i>Rhantus frontalis</i> (Marsham, 1802)	<b>B4</b>	
	<i>Rhantus grapii</i> (Gyllenhal, 1808)	<b>B4</b>	
	<i>Rhantus suturalis</i> (MacLeay, 1825)	<b>B4</b>	
	<i>Rhantus suturellus</i> (Harris, 1828)	<b>B4</b>	
	<i>Rhantus</i> sp.		<b>65</b>
(Dytiscinae)	<i>Acilius canaliculatus</i> (Nicolai, 1822)	<b>B6</b>	
	<i>Acilius sulcatus</i> (Linnaeus, 1758)	<b>B6</b>	
	<i>Acilius</i> sp.		<b>30</b>
	<i>Cybister lateralimarginalis</i> (De Geer, 1774)	<b>B6</b>	<b>31</b>
	<i>Dytiscus circumcinctus</i> Ahrens, 1811	<b>B6</b>	
	<i>Dytiscus circumflexus</i> Fabricius, 1801	<b>B6</b>	
	<i>Dytiscus dimidiatus</i> Bergstrasser, 1778	<b>B6</b>	
	<i>Dytiscus latissimus</i> Linnaeus, 1758	<b>B6</b>	
	<i>Dytiscus marginalis</i> Linnaeus, 1758	<b>B6</b>	
	<i>Dytiscus semisulcatus</i> O.F. Müller, 1776	<b>B6</b>	
	<i>Dytiscus</i> sp.		<b>32</b>
	<i>Graphoderus austriacus</i> (Sturm, 1834)	<b>B6</b>	
	<i>Graphoderus cinereus</i> (Linnaeus, 1758)	<b>B6</b>	
	<i>Graphoderus</i> sp.		<b>33</b>
	<i>Hydaticus aruspex</i> Clark, 1864	<b>B6</b>	
	<i>Hydaticus continentalis</i> J. Balfour-Browne, 1944	<b>B6</b>	
	<i>Hydaticus seminiger</i> (De Geer, 1774)	<b>B6</b>	
	<i>Hydaticus transversalis</i> (Pontoppidan, 1763)	<b>B6</b>	
	<i>Hydaticus</i> sp.		<b>34</b>
<b>Helophoridae</b>	<i>Helophorus aequalis</i> Thomson, 1868	<b>B7</b>	
	<i>Helophorus aquaticus</i> (Linnaeus, 1758)	<b>B7</b>	
	<i>Helophorus arvernicus</i> Mulsant, 1846	<b>B7</b>	
	<i>Helophorus asperatus</i> Rey, 1885	<b>B7</b>	
	<i>Helophorus brevipalpis</i> Bedel, 1881	<b>B7</b>	
	<i>Helophorus discrepans</i> Rey, 1885	<b>B7</b>	
	<i>Helophorus flavipes</i> Fabricius, 1792	<b>B7</b>	
	<i>Helophorus glacialis</i> Villa, 1883	<b>B7</b>	
	<i>Helophorus grandis</i> Illiger, 1798	<b>B7</b>	
	<i>Helophorus granularis</i> (Linnaeus, 1761)	<b>B7</b>	
	<i>Helophorus griseus</i> Herbst, 1793	<b>B7</b>	
	<i>Helophorus minutus</i> Fabricius, 1775	<b>B7</b>	
	<i>Helophorus montenegrinus</i> Kuwert, 1885	<b>B7</b>	
	<i>Helophorus nanus</i> Sturm, 1836	<b>B7</b>	
	<i>Helophorus redtenbacheri</i> Kuwert, 1885	<b>B7</b>	
<b>Georissidae</b>	<i>Georissus crenulatus</i> (Rossi, 1794)	<b>B8</b>	
<b>Hydrochidae</b>	<i>Hydrochus crenatus</i> (Fabricius, 1792)	<b>B8</b>	
	<i>Hydrochus elongatus</i> (Schaller, 1783)	<b>B8</b>	
	<i>Hydrochus ignicollis</i> Motschulsky, 1860	<b>B8</b>	
	<i>Hydrochus megaphallus</i> Berge Henegouwen, 1988	<b>B8</b>	
<b>Spercheidae</b>	<i>Spercheus emarginatus</i> (Schaller, 1783)	<b>B8</b>	<b>35</b>
<b>Hydrophilidae</b>			
(Sphaeridiinae)	<i>Cercyon bifenestratus</i> Küster, 1851	<b>B8</b>	
	<i>Coelostoma orbiculare</i> (Fabricius, 1775)	<b>B8</b>	<b>36</b>
(Hydrophilinae)	<i>Anacaena bipustulata</i> (Marsham, 1802)	<b>B9</b>	
	<i>Anacaena globulus</i> (Paykull, 1798)	<b>B9</b>	
	<i>Anacaena limbata</i> (Fabricius, 1792)	<b>B9</b>	
	<i>Anacaena lutescens</i> (Stephens, 1829)	<b>B9</b>	
	<i>Anacaena</i> sp.		<b>37</b>
	<i>Berosus frontifoveatus</i> Kuwert, 1888	<b>B9</b>	

	<i>Berosus signaticollis</i> (Charpentier, 1825)	<b>B9</b>	
	<i>Berosus spinosus</i> (Steven, 1808)	<b>B9</b>	
	<i>Berosus</i> sp.		<b>66</b>
	<i>Crenitis punctatostrata</i> (Letzner, 1840)	<b>B9</b>	
	<i>Cymbiodyta marginella</i> (Fabricius, 1792)	<b>B9</b>	<b>38</b>
	<i>Enochrus affinis</i> (Thunberg, 1794)	<b>B10</b>	
	<i>Enochrus bicolor</i> (Fabricius, 1792)	<b>B10</b>	
	<i>Enochrus coarctatus</i> (Gredler, 1863)	<b>B10</b>	
	<i>Enochrus melanocephalus</i> (Olivier, 1792)	<b>B10</b>	
	<i>Enochrus ochropterus</i> (Marsham, 1802)	<b>B10</b>	<b>67</b>
	<i>Enochrus quadripunctatus</i> (Herbst, 1797)	<b>B10</b>	
	<i>Enochrus testaceus</i> (Fabricius, 1801)	<b>B10</b>	
	<i>Enochrus</i> sp.		<b>39</b>
	<i>Helochares lividus</i> (Forster, 1771)	<b>B10</b>	
	<i>Helochares obscurus</i> (Müller, 1776)	<b>B10</b>	
	<i>Helochares</i> sp.		<b>68</b>
	<i>Hydrobius fuscipes</i> (Linnaeus, 1758)	<b>B10</b>	<b>40</b>
	<i>Hydrochara caraboides</i> (Linnaeus, 1758)	<b>B10</b>	
	<i>Hydrochara flavipes</i> (Steven, 1808)	<b>B10</b>	
	<i>Hydrochara</i> sp.		<b>41</b>
	<i>Hydrophilus atterimus</i> (Eschscholtz, 1822)	<b>B10</b>	
	<i>Hydrophilus piceus</i> (Linnaeus, 1758)	<b>B10</b>	
	<i>Hydrophilus</i> sp.		<b>42</b>
	<i>Chaetarthria seminulum</i> (Herbst, 1797)	<b>B9</b>	
	<i>Laccobius atratus</i> Rottenberg, 1874	<b>B9</b>	
	<i>Laccobius bipunctatus</i> (Fabricius, 1775)	<b>B9</b>	
	<i>Laccobius minutus</i> (Linnaeus, 1758)	<b>B9</b>	
	<i>Laccobius striatulus</i> (Fabricius, 1801)	<b>B9</b>	
	<i>Laccobius</i> sp.		<b>43</b>
	<i>Limnoxenus niger</i> (Zschach, 1788)	<b>B10</b>	
	<i>Paracymus aeneus</i> (Germar, 1824)	<b>B9</b>	
<b>Hydraenidae</b>	<i>Hydraena belgica</i> d'Orchymont, 1930	<b>B11</b>	
	<i>Hydraena britteni</i> Joy, 1907	<b>B11</b>	
	<i>Hydraena dentipes</i> Germar, 1844	<b>B11</b>	
	<i>Hydraena excisa</i> Kiesenwetter, 1849	<b>B11</b>	
	<i>Hydraena gracilis</i> Germar, 1824	<b>B11</b>	
	<i>Hydraena lapidicola</i> Kiesenwetter, 1849	<b>B11</b>	
	<i>Hydraena melas</i> Dalla Torre, 1877	<b>B11</b>	
	<i>Hydraena minutissima</i> Stephens, 1829	<b>B11</b>	
	<i>Hydraena morio</i> Kiesenwetter, 1849	<b>B11</b>	
	<i>Hydraena nigrita</i> Germar, 1824	<b>B11</b>	
	<i>Hydraena paganettii</i> Ganglbauer, 1901	<b>B11</b>	
	<i>Hydraena palustris</i> Erichson, 1837	<b>B11</b>	
	<i>Hydraena pulchella</i> Germar, 1824	<b>B11</b>	
	<i>Hydraena pygmaea</i> Waterhouse, 1833	<b>B11</b>	
	<i>Hydraena reyi</i> Kuwert, 1888	<b>B11</b>	
	<i>Hydraena riparia</i> Kugelann, 1794	<b>B11</b>	
	<i>Hydraena saga</i> D'orchymont, 1930	<b>B11</b>	
	<i>Hydraena schuleri</i> Ganglbauer, 1901	<b>B11</b>	
	<i>Hydraena truncata</i> Rey, 1885	<b>B11</b>	
	<i>Limnebius aluta</i> Bedel, 1881	<b>B12</b>	
	<i>Limnebius papposus</i> Mulsant, 1844	<b>B12</b>	
	<i>Limnebius truncatellus</i> (Thunberg, 1794)	<b>B12</b>	
	<i>Ochthebius melanescens</i> Dalla Torre, 1877	<b>B12</b>	
	<i>Ochthebius metallescens</i> Rosenhauer, 1847	<b>B12</b>	
	<i>Ochthebius minimus</i> (Fabricius, 1792)	<b>B12</b>	

	<i>Ochthebius steinbuehleri</i> (Reitter, 1886)		44
<b>Scirtidae</b>	<i>Cyphon phragmiteticola</i> Nyholm, 1955	<b>B13</b>	
	<i>Cyphon</i> spp.		69
	<i>Elodes marginata</i> (Fabricius, 1798)	<b>B13</b>	70
	<i>Elodes</i> gr. <i>minuta</i>		45
	<i>Hydrocyphon deflexicollis</i> (P. W. J. Müller 1821)		46
	<i>Microcara testacea</i> (Linnaeus, 1767)	<b>B13</b>	47
	<i>Scirtes hemisphaericus</i> (Linnaeus, 1767)	<b>B13</b>	
	<i>Scirtes</i> sp.		48
<b>Psephenidae</b>	<i>Eubria palustris</i> (Germar, 1818)	<b>B8</b>	49
<b>Dryopidae</b>	<i>Dryops auriculatus</i> (Geoffroy, 1785)	<b>B13</b>	
<b>Elimidae</b>	<i>Elmis aenea</i> (Müller, 1806)	<b>B13</b>	50
	<i>Elmis latreillei</i> Bedel, 1878	<b>B13</b>	51
	<i>Elmis maugetii maugetii</i> Latreille, 1798	<b>B13</b>	52
	<i>Elmis obscura</i> (Müller, 1806)	<b>B13</b>	
	<i>Elmis rioloides</i> Kuwert, 1890	<b>B13</b>	
	<i>Esolus angustatus</i> (Müller, 1821)	<b>B13</b>	
	<i>Esolus parallelepipedus</i> (Müller, 1806)	<b>B13</b>	
	<i>Esolus</i> sp.		53
	<i>Limnius perrisi perrisi</i> (Dufour, 1843)	<b>B13</b>	54
	<i>Limnius volckmari</i> (Panzer, 1793)	<b>B13</b>	55
	<i>Macronychus quadrituberculatus</i> Müller, 1806	<b>B13</b>	56
	<i>Oulimnius tuberculatus</i> (Müller, 1806)	<b>B13</b>	57
	<i>Riolus subviolaceus</i> (Müller, 1817)	<b>B13</b>	58
	<i>Potamophilus acuminatus</i> (Fabricius, 1792)	<b>B13</b>	59
<b>Limnichidae</b>	<i>Pelochares versicolor</i> (Waltl, 1833)	<b>B13</b>	
<b>Curculionidae</b>	<i>Tanysphyrus lemnae</i> (Paykull, 1792)	<b>B8</b>	

**determinovali:** Mgr. Michal Straka, Mgr. Jan Sychra a další  
**místo uložení:** ÚBZ, budova č. 10, pracovna hydrobiologie - dveře č. 222; tel. 532146342  
**kontaktní osoba:** Mgr. Jan Sychra (dubovec@seznam.cz)